

# MD7513 - Prevention, Screening & Early Detection in Diabetes

Prevention, Screening & Early Detection in Diabetes

View Online



[1]

A community based primary prevention programme for type 2 diabetes integrating identification and lifestyle intervention for prevention: the Let's Prevent Diabetes cluster randomised controlled trial- ClinicalKey:  
<https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S0091743515003850>.

[2]

Absetz, P. et al. 2009. Type 2 Diabetes Prevention in the Real World: Three-year results of the GOAL Lifestyle Implementation Trial. *Diabetes Care*. 32, 8 (Aug. 2009), 1418-1420.  
DOI:<https://doi.org/10.2337/dc09-0039>.

[3]

Ackermann, R.T. et al. 2008. Translating the Diabetes Prevention Program into the Community. *American Journal of Preventive Medicine*. 35, 4 (Oct. 2008), 357-363.  
DOI:<https://doi.org/10.1016/j.amepre.2008.06.035>.

[4]

Aguiar, E.J. et al. 2016. Efficacy of the Type 2 Diabetes Prevention Using LifeStyle Education Program RCT. *American Journal of Preventive Medicine*. 50, 3 (Mar. 2016), 353-364. DOI:<https://doi.org/10.1016/j.amepre.2015.08.020>.

[5]

Aune, D. et al. 2013. Whole grain and refined grain consumption and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies.

European Journal of Epidemiology. 28, 11 (Nov. 2013), 845–858.  
DOI:<https://doi.org/10.1007/s10654-013-9852-5>.

[6]

Bailey, Stephen Academic Writing : A Handbook for International Students.

[7]

Barry, E. et al. 2017. Efficacy and effectiveness of screen and treat policies in prevention of type 2 diabetes: systematic review and meta-analysis of screening tests and interventions. *BMJ*. (Jan. 2017). DOI:<https://doi.org/10.1136/bmj.i6538>.

[8]

Breeze, P.R. et al. 2017. Cost-effectiveness of population-based, community, workplace and individual policies for diabetes prevention in the UK. *Diabetic Medicine*. 34, 8 (Aug. 2017), 1136–1144. DOI:<https://doi.org/10.1111/dme.13349>.

[9]

Breeze, P.R. et al. 2017. The impact of Type 2 diabetes prevention programmes based on risk-identification and lifestyle intervention intensity strategies: a cost-effectiveness analysis. *Diabetic Medicine*. 34, 5 (May 2017), 632–640.  
DOI:<https://doi.org/10.1111/dme.13314>.

[10]

Brown, J. et al. 1996. Lifestyle interventions for the treatment of women with gestational diabetes. *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd.

[11]

Carlsson, L.M.S. et al. 2012. Bariatric Surgery and Prevention of Type 2 Diabetes in Swedish Obese Subjects. *New England Journal of Medicine*. 367, 8 (Aug. 2012), 695–704.  
DOI:<https://doi.org/10.1056/NEJMoa1112082>.

[12]

Carlsson, L.M.S. et al. 2012. Bariatric Surgery and Prevention of Type 2 Diabetes in Swedish Obese Subjects. *New England Journal of Medicine*. 367, 8 (Aug. 2012), 695–704. DOI:<https://doi.org/10.1056/NEJMoa1112082>.

[13]

Carter, P. et al. 2012. Dietary Recommendations for the Prevention of Type 2 diabetes: What Are They Based on? *Journal of Nutrition and Metabolism*. 2012, (2012), 1–6. DOI:<https://doi.org/10.1155/2012/847202>.

[14]

Church, T.S. et al. 2010. Effects of Aerobic and Resistance Training on Hemoglobin A Levels in Patients With Type 2 Diabetes. *JAMA*. 304, 20 (Nov. 2010). DOI:<https://doi.org/10.1001/jama.2010.1710>.

[15]

Clinical applicability and cost-effectiveness of DIABSCORE in screening for type 2 diabetes in primary care- ClinicalKey:  
<https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S0168822717301031>.

[16]

Colberg, S.R. et al. 2010. Exercise and Type 2 Diabetes: The American College of Sports Medicine and the American Diabetes Association: joint position statement executive summary. *Diabetes Care*. 33, 12 (Dec. 2010), 2692–2696. DOI:<https://doi.org/10.2337/dc10-1548>.

[17]

Colberg, S.R. et al. 2016. Physical Activity/Exercise and Diabetes: A Position Statement of the American Diabetes Association. *Diabetes Care*. 39, 11 (Nov. 2016), 2065–2079. DOI:<https://doi.org/10.2337/dc16-1728>.

[18]

Creame, P. and Lea, M.R. 2008. Writing At University: A Guide For Students. McGraw-Hill Education.

[19]

Dagfinn Aune 2016. Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies. The BMJ. 353, (2016). DOI:<https://doi.org/doi:10.1136/bmj.i2716>.

[20]

Deborah F Tate 2012. Replacing caloric beverages with water or diet beverages for weight loss in adults: main results of the Choose Healthy Options Consciously Everyday (CHOICE) randomized clinical trial. The American Journal of Clinical Nutrition. 95, 3 (2012). DOI:<https://doi.org/doi:10.3945/ajcn.111.026278>.

[21]

Dhingra, R. et al. 2007. Soft Drink Consumption and Risk of Developing Cardiometabolic Risk Factors and the Metabolic Syndrome in Middle-Aged Adults in the Community. Circulation. 116, 5 (Jul. 2007), 480–488. DOI:<https://doi.org/10.1161/CIRCULATIONAHA.107.689935>.

[22]

Differences in the prospective association between individual plasma phospholipid saturated fatty acids and incident type 2 diabetes: the EPIC-InterAct case-cohort study: [http://ac.els-cdn.com/S2213858714701469/1-s2.0-S2213858714701469-main.pdf?\\_tid=41970bbc-7e93-11e7-a8e5-00000aab0f01&acdnat=1502455838\\_1e4678aef6e5b79c43c42f6b68f370f1](http://ac.els-cdn.com/S2213858714701469/1-s2.0-S2213858714701469-main.pdf?_tid=41970bbc-7e93-11e7-a8e5-00000aab0f01&acdnat=1502455838_1e4678aef6e5b79c43c42f6b68f370f1).

[23]

Dyson, P.A. et al. 2011. Diabetes UK evidence-based nutrition guidelines for the prevention and management of diabetes. Diabetic Medicine. 28, 11 (Nov. 2011), 1282–1288. DOI:<https://doi.org/10.1111/j.1464-5491.2011.03371.x>.

[24]

Effect of a lifestyle intervention on weight change in south Asian individuals in the UK at high risk of type 2 diabetes: a family-cluster randomised controlled trial- ClinicalKey: <https://www.clinicalkey.com/#!/content/journal/1-s2.0-S2213858713702043>.

[25]

Emily A Hu 2012. White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review. *The BMJ*. 344, (2012). DOI:<https://doi.org/doi:10.1136/bmj.e1454>.

[26]

Fogelholm, M. et al. 2017. PREVIEW: Prevention of Diabetes through Lifestyle Intervention and Population Studies in Europe and around the World. Design, Methods, and Baseline Participant Description of an Adult Cohort Enrolled into a Three-Year Randomised Clinical Trial. *Nutrients*. 9, 6 (Jun. 2017). DOI:<https://doi.org/10.3390/nu9060632>.

[27]

Georgios S Papaetis 2014. Incretin-based therapies in prediabetes: Current evidence and future perspectives. *World Journal of Diabetes*. 5, 6 (2014). DOI:<https://doi.org/doi:10.4239/wjd.v5.i6.817>.

[28]

Gillies, C.L. et al. 2007. Pharmacological and lifestyle interventions to prevent or delay type 2 diabetes in people with impaired glucose tolerance: systematic review and meta-analysis. *BMJ*. 334, 7588 (Jan. 2007), 299-299. DOI:<https://doi.org/10.1136/bmj.39063.689375.55>.

[29]

Gopalan, A. et al. 2015. Awareness of Prediabetes and Engagement in Diabetes Risk-Reducing Behaviors. *American Journal of Preventive Medicine*. 49, 4 (Oct. 2015), 512-519. DOI:<https://doi.org/10.1016/j.amepre.2015.03.007>.

[30]

Gray, L.J. et al. 2012. Detection of impaired glucose regulation and/or type 2 diabetes mellitus, using primary care electronic data, in a multiethnic UK community setting.

Diabetologia. 55, 4 (Apr. 2012), 959–966. DOI:<https://doi.org/10.1007/s00125-011-2432-x>.

[31]

Gray, L.J. et al. 2016. Engagement, Retention, and Progression to Type 2 Diabetes: A Retrospective Analysis of the Cluster-Randomised 'Let's Prevent Diabetes' Trial. *PLOS Medicine*. 13, 7 (Jul. 2016). DOI:<https://doi.org/10.1371/journal.pmed.1002078>.

[32]

Gray, L.J. et al. 2010. The Leicester Risk Assessment score for detecting undiagnosed Type 2 diabetes and impaired glucose regulation for use in a multiethnic UK setting. *Diabetic Medicine*. 27, 8 (May 2010), 887–895.  
DOI:<https://doi.org/10.1111/j.1464-5491.2010.03037.x>.

[33]

Gregg, E.W. and Shaw, J.E. 2017. Global Health Effects of Overweight and Obesity. *New England Journal of Medicine*. (Jun. 2017). DOI:<https://doi.org/10.1056/NEJMe1706095>.

[34]

Hamman, R.F. et al. 2006. Effect of Weight Loss With Lifestyle Intervention on Risk of Diabetes. *Diabetes Care*. 29, 9 (Sep. 2006), 2102–2107.  
DOI:<https://doi.org/10.2337/dc06-0560>.

[35]

Helen C Eborall 2007. Psychological impact of screening for type 2 diabetes: controlled trial and comparative study embedded in the ADDITION (Cambridge) randomised controlled trial. *BMJ : British Medical Journal*. 335, 7618 (2007). DOI:<https://doi.org/doi:10.1136/bmj.39303.723449.55>.

[36]

How to prevent type 2 diabetes in women with previous gestational diabetes? A systematic review of behavioural interventions- ClinicalKey:  
<https://www.clinicalkey.com/#!/content/playContent/1-s2.0-S1751991817300785?returnurl>

=null&referrer=null.

[37]

Joiner, K.L. et al. 2017. Lifestyle interventions based on the diabetes prevention program delivered via eHealth: A systematic review and meta-analysis. *Preventive Medicine*. 100, (Jul. 2017), 194–207. DOI:<https://doi.org/10.1016/j.ypmed.2017.04.033>.

[38]

Koivusalo, S.B. et al. 2016. Gestational Diabetes Mellitus Can Be Prevented by Lifestyle Intervention: The Finnish Gestational Diabetes Prevention Study (RADIEL). *Diabetes Care*. 39, 1 (Jan. 2016), 24–30. DOI:<https://doi.org/10.2337/dc15-0511>.

[39]

Laaksonen, D.E. et al. 2005. Physical Activity in the Prevention of Type 2 Diabetes: The Finnish Diabetes Prevention Study. *Diabetes*. 54, 1 (Jan. 2005), 158–165. DOI:<https://doi.org/10.2337/diabetes.54.1.158>.

[40]

LaMonte, M.J. 2005. Physical activity and diabetes prevention. *Journal of Applied Physiology*. 99, 3 (Sep. 2005), 1205–1213. DOI:<https://doi.org/10.1152/jappphysiol.00193.2005>.

[41]

Less Sitting, More Physical Activity, or Higher Fitness?- ClinicalKey:  
<https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/playContent/1-s2.0-S0025619615006308?returnurl=null&referrer=null>.

[42]

Ley, S.H. et al. 2014. Prevention and management of type 2 diabetes: dietary components and nutritional strategies. *The Lancet*. 383, 9933 (Jun. 2014), 1999–2007. DOI:[https://doi.org/10.1016/S0140-6736\(14\)60613-9](https://doi.org/10.1016/S0140-6736(14)60613-9).

[43]

Li, R. et al. 2010. Cost-Effectiveness of Interventions to Prevent and Control Diabetes Mellitus: A Systematic Review. *Diabetes Care*. 33, 8 (Aug. 2010), 1872–1894.  
DOI:<https://doi.org/10.2337/dc10-0843>.

[44]

Linda Penn 2013. Importance of Weight Loss Maintenance and Risk Prediction in the Prevention of Type 2 Diabetes: Analysis of European Diabetes Prevention Study RCT. *PLoS ONE*. 8, 2 (2013). DOI:<https://doi.org/doi:10.1371/journal.pone.0057143>.

[45]

Linda Penn 2013. Importance of Weight Loss Maintenance and Risk Prediction in the Prevention of Type 2 Diabetes: Analysis of European Diabetes Prevention Study RCT. *PLoS ONE*. 8, 2 (2013). DOI:<https://doi.org/doi:10.1371/journal.pone.0057143>.

[46]

Lindstrom, J. et al. 2003. The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care*. 26, 12 (Dec. 2003), 3230–3236. DOI:<https://doi.org/10.2337/diacare.26.12.3230>.

[47]

Long, G.H. et al. 2015. Healthy behaviours and 10-year incidence of diabetes: A population cohort study. *Preventive Medicine*. 71, (Feb. 2015), 121–127.  
DOI:<https://doi.org/10.1016/j.ypmed.2014.12.013>.

[48]

Luchsinger, J.A. et al. 2017. Metformin, Lifestyle Intervention, and Cognition in the Diabetes Prevention Program Outcomes Study. *Diabetes Care*. 40, 7 (Jul. 2017), 958–965.  
DOI:<https://doi.org/10.2337/dc16-2376>.

[49]

Malik, V.S. et al. 2010. Sugar-Sweetened Beverages and Risk of Metabolic Syndrome and



Type 2 Diabetes: A meta-analysis. *Diabetes Care*. 33, 11 (Nov. 2010), 2477–2483.  
DOI:<https://doi.org/10.2337/dc10-1079>.

[50]

Malik, V.S. et al. 2013. Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *American Journal of Clinical Nutrition*. 98, 4 (Oct. 2013), 1084–1102. DOI:<https://doi.org/10.3945/ajcn.113.058362>.

[51]

Mann, J. et al. 2016. Dietary guidelines on trial: the charges are not evidence based. *The Lancet*. 388, 10047 (Aug. 2016), 851–853.  
DOI:[https://doi.org/10.1016/S0140-6736\(16\)31278-8](https://doi.org/10.1016/S0140-6736(16)31278-8).

[52]

Matti Uusitupa 2009. Ten-Year Mortality and Cardiovascular Morbidity in the Finnish Diabetes Prevention Study—Secondary Analysis of the Randomized Trial. *PLoS ONE*. 4, 5 (2009). DOI:<https://doi.org/doi:10.1371/journal.pone.0005656>.

[53]

Nathan, D.M. et al. 2007. Impaired Fasting Glucose and Impaired Glucose Tolerance: Implications for care. *Diabetes Care*. 30, 3 (Mar. 2007), 753–759.  
DOI:<https://doi.org/10.2337/dc07-9920>.

[54]

Orozco, L.J. et al. 1996. Exercise or exercise and diet for preventing type 2 diabetes mellitus. *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd.

[55]

Pan, A. et al. 2011. Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *American Journal of Clinical Nutrition*. 94, 4 (Oct. 2011), 1088–1096. DOI:<https://doi.org/10.3945/ajcn.111.018978>.

[56]

Pan, X.-R. et al. 1997. Effects of Diet and Exercise in Preventing NIDDM in People With Impaired Glucose Tolerance: The Da Qing IGT and Diabetes Study. *Diabetes Care*. 20, 4 (Apr. 1997), 537-544. DOI:<https://doi.org/10.2337/diacare.20.4.537>.

[57]

Rajul Parikh, Annie Mathai, Shefali Parikh, G. Chandra Sekhar, Ravi Thomas Understanding and using sensitivity, specificity and predictive values. *Indian Journal of Ophthalmology*.

[58]

Ramachandran, A. et al. 2006. The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia*. 49, 2 (Feb. 2006), 289-297. DOI:<https://doi.org/10.1007/s00125-005-0097-z>.

[59]

Schulze, M.B. 2004. Sugar-Sweetened Beverages, Weight Gain, and Incidence of Type 2 Diabetes in Young and Middle-Aged Women. *JAMA*. 292, 8 (Aug. 2004). DOI:<https://doi.org/10.1001/jama.292.8.927>.

[60]

Schwingshackl, L. et al. 2017. Food groups and risk of type 2 diabetes mellitus: a systematic review and meta-analysis of prospective studies. *European Journal of Epidemiology*. 32, 5 (May 2017), 363-375. DOI:<https://doi.org/10.1007/s10654-017-0246-y>.

[61]

Sedentary Behaviors and Subsequent Health Outcomes in Adults- ClinicalKey:  
<https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/journal/1-s2.0-S0749379711003126>.

[62]

Sedentary Behaviors and Subsequent Health Outcomes in Adults- ClinicalKey:  
<https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/journal/1-s2.0-S0749379711003126>.

[63]

Stefan, N. et al. 2015. A high-risk phenotype associates with reduced improvement in glycaemia during a lifestyle intervention in prediabetes. *Diabetologia*. 58, 12 (Dec. 2015), 2877–2884. DOI:<https://doi.org/10.1007/s00125-015-3760-z>.

[64]

Steven N Blair 2009. Physical inactivity: the biggest public health problem of the 21st century. *British Journal of Sports Medicine*. 43, (Jan. 2009), 1–2.

[65]

Sweeting, A.N. et al. 2016. Gestational Diabetes Mellitus in Early Pregnancy: Evidence for Poor Pregnancy Outcomes Despite Treatment. *Diabetes Care*. 39, 1 (Jan. 2016), 75–81. DOI:<https://doi.org/10.2337/dc15-0433>.

[66]

Thomas Yates 2012. Walking away from type 2 diabetes: trial protocol of a cluster randomised controlled trial evaluating a structured education programme in those at high risk of developing type 2 diabetes. *BMC Family Practice*. 13, (2012). DOI:<https://doi.org/doi:10.1186/1471-2296-13-46>.

[67]

Tseng, E. et al. 2017. Metformin Use in Prediabetes Among U.S. Adults, 2005–2012. *Diabetes Care*. 40, 7 (Jul. 2017), 887–893. DOI:<https://doi.org/10.2337/dc16-1509>.

[68]

Tuomilehto, J. et al. 2001. Prevention of Type 2 Diabetes Mellitus by Changes in Lifestyle among Subjects with Impaired Glucose Tolerance. *New England Journal of Medicine*. 344, 18 (May 2001), 1343–1350. DOI:<https://doi.org/10.1056/NEJM200105033441801>.

[69]

Vita, P. et al. 2016. Type 2 diabetes prevention in the community: 12-Month outcomes from the Sydney Diabetes Prevention Program. *Diabetes Research and Clinical Practice*. 112, (Feb. 2016), 13-19. DOI:<https://doi.org/10.1016/j.diabres.2015.11.010>.

[70]

Webb, D.R. et al. 2011. Screening for diabetes using an oral glucose tolerance test within a western multi-ethnic population identifies modifiable cardiovascular risk: the ADDITION-Leicester study. *Diabetologia*. 54, 9 (Sep. 2011), 2237-2246. DOI:<https://doi.org/10.1007/s00125-011-2189-2>.

[71]

Wu, Y. et al. 2015. Fruit and vegetable consumption and risk of type 2 diabetes mellitus: A dose-response meta-analysis of prospective cohort studies. *Nutrition, Metabolism and Cardiovascular Diseases*. 25, 2 (Feb. 2015), 140-147. DOI:<https://doi.org/10.1016/j.numecd.2014.10.004>.

[72]

Yates, T. et al. 2011. Stand up for your health: Is it time to rethink the physical activity paradigm? *Diabetes Research and Clinical Practice*. 93, 2 (Aug. 2011), 292-294. DOI:<https://doi.org/10.1016/j.diabres.2011.03.023>.

[73]

20AD. A Systems Science Approach to the Obesity Epidemic and Prevention of Diabetes.

[74]

2017. American Diabetes Association - Standards of Medical Care in Diabetes - 2017 Summary of Revisions. *Diabetes Care*. 40, Supplement 1 (Jan. 2017), S4-S5. DOI:<https://doi.org/10.2337/dc17-S003>.

[75]

2007. Correction. *Circulation*. 116, 23 (Dec. 2007), e557–e557.  
DOI:<https://doi.org/10.1161/CIRCULATIONAHA.107.187928>.

[76]

2017. Health Effects of Overweight and Obesity in 195 Countries over 25 Years. *New England Journal of Medicine*. (Jun. 2017). DOI:<https://doi.org/10.1056/NEJMoa1614362>.

[77]

NICE - Type 2 diabetes: prevention in people at high risk | Guidance and guidelines. NICE.

[78]

2008. Nutrition Recommendations and Interventions for Diabetes: A position statement of the American Diabetes Association. *Diabetes Care*. 31, Supplement 1 (Jan. 2008), S61–S78.  
DOI:<https://doi.org/10.2337/dc08-S061>.

[79]

Obesity: identification, assessment and management | Guidance and guidelines | NICE.

[80]

2015. Public Health England - Diabetes prevention programmes: evidence review. Public Health England.

[81]

2002. Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin. *New England Journal of Medicine*. 346, 6 (Feb. 2002), 393–403.  
DOI:<https://doi.org/10.1056/NEJMoa012512>.

[82]

2002. Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin. *New England Journal of Medicine*. 346, 6 (Feb. 2002), 393–403.  
DOI:<https://doi.org/10.1056/NEJMoa012512>.

[83]

2007. Screening for type 2 diabetes: literature review and economic modelling. *Clinical Governance: An International Journal*. 12, 4 (Oct. 2007).  
DOI:<https://doi.org/10.1108/cgij.2007.24812dae.003>.

[84]

2012. The 10-Year Cost-Effectiveness of Lifestyle Intervention or Metformin for Diabetes Prevention: An intent-to-treat analysis of the DPP/DPPOS. *Diabetes Care*. 35, 4 (Apr. 2012), 723–730. DOI:<https://doi.org/10.2337/dc11-1468>.

[85]

Type 2 diabetes: prevention in people at high risk | Guidance and guidelines | NICE.