

MD7513 - Prevention, Screening & Early Detection in Diabetes

Prevention, Screening & Early Detection in Diabetes

View Online



A community based primary prevention programme for type 2 diabetes integrating identification and lifestyle intervention for prevention: the Let's Prevent Diabetes cluster randomised controlled trial- ClinicalKey. (n.d.).
<https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S0091743515003850>

A Systems Science Approach to the Obesity Epidemic and Prevention of Diabetes. (20 C.E.).
<https://www.youtube.com/watch?v=HrrD-Z0B7dY&feature=youtu.be&t=55s%20A%20Systems%20Science%20Approach%20to%20the%20Obesity%20Epidemic%20and%20Prevention%20of%20Diabetes>.

Absetz, P., Oldenburg, B., Hankonen, N., Valve, R., Heinonen, H., Nissinen, A., Fogelholm, M., Talja, M., & Uutela, A. (2009). Type 2 Diabetes Prevention in the Real World: Three-year results of the GOAL Lifestyle Implementation Trial. *Diabetes Care*, 32(8), 1418–1420. <https://doi.org/10.2337/dc09-0039>

Ackermann, R. T., Finch, E. A., Brizendine, E., Zhou, H., & Marrero, D. G. (2008). Translating the Diabetes Prevention Program into the Community. *American Journal of Preventive Medicine*, 35(4), 357–363. <https://doi.org/10.1016/j.amepre.2008.06.035>

Aguiar, E. J., Morgan, P. J., Collins, C. E., Plotnikoff, R. C., Young, M. D., & Callister, R. (2016). Efficacy of the Type 2 Diabetes Prevention Using LifeStyle Education Program RCT. *American Journal of Preventive Medicine*, 50(3), 353–364. <https://doi.org/10.1016/j.amepre.2015.08.020>

American Diabetes Association - Standards of Medical Care in Diabetes - 2017 Summary of Revisions. (2017). *Diabetes Care*, 40(Supplement 1), S4–S5. <https://doi.org/10.2337/dc17-S003>

Aune, D., Norat, T., Romundstad, P., & Vatten, L. J. (2013). Whole grain and refined grain consumption and the risk of type 2 diabetes: a systematic review and dose–response meta-analysis of cohort studies. *European Journal of Epidemiology*, 28(11), 845–858. <https://doi.org/10.1007/s10654-013-9852-5>

Bailey, Stephen. (n.d.). *Academic Writing: A Handbook for International Students* (5th ed.). <https://ebookcentral.proquest.com/lib/leicester/detail.action?docID=5178437>

Barry, E., Roberts, S., Oke, J., Vijayaraghavan, S., Normansell, R., & Greenhalgh, T. (2017). Efficacy and effectiveness of screen and treat policies in prevention of type 2 diabetes:

systematic review and meta-analysis of screening tests and interventions. *BMJ*.
<https://doi.org/10.1136/bmj.i6538>

Breeze, P. R., Thomas, C., Squires, H., Brennan, A., Greaves, C., Diggle, P., Brunner, E., Tabak, A., Preston, L., & Chilcott, J. (2017a). Cost-effectiveness of population-based, community, workplace and individual policies for diabetes prevention in the UK. *Diabetic Medicine*, 34(8), 1136–1144. <https://doi.org/10.1111/dme.13349>

Breeze, P. R., Thomas, C., Squires, H., Brennan, A., Greaves, C., Diggle, P. J., Brunner, E., Tabak, A., Preston, L., & Chilcott, J. (2017b). The impact of Type 2 diabetes prevention programmes based on risk-identification and lifestyle intervention intensity strategies: a cost-effectiveness analysis. *Diabetic Medicine*, 34(5), 632–640.
<https://doi.org/10.1111/dme.13314>

Brown, J., Alwan, N. A., West, J., Brown, S., McKinlay, C. J., Farrar, D., & Crowther, C. A. (1996). Lifestyle interventions for the treatment of women with gestational diabetes. In *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd.
<https://doi.org/10.1002/14651858.CD011970.pub2>

Carlsson, L. M. S., Peltonen, M., Ahlin, S., Anveden, A., Bouchard, C., Carlsson, B., Jacobson, P., Lonroth, H., Maglio, C., Naslund, I., Pirazzi, C., Romeo, S., Sjöholm, K., Sjöström, E., Wedel, H., Svensson, P.-A., & Sjöström, L. (2012a). Bariatric Surgery and Prevention of Type 2 Diabetes in Swedish Obese Subjects. *New England Journal of Medicine*, 367(8), 695–704. <https://doi.org/10.1056/NEJMoa1112082>

Carlsson, L. M. S., Peltonen, M., Ahlin, S., Anveden, A., Bouchard, C., Carlsson, B., Jacobson, P., Lonroth, H., Maglio, C., Naslund, I., Pirazzi, C., Romeo, S., Sjöholm, K., Sjöström, E., Wedel, H., Svensson, P.-A., & Sjöström, L. (2012b). Bariatric Surgery and Prevention of Type 2 Diabetes in Swedish Obese Subjects. *New England Journal of Medicine*, 367(8), 695–704. <https://doi.org/10.1056/NEJMoa1112082>

Carter, P., Khunti, K., & Davies, M. J. (2012). Dietary Recommendations for the Prevention of Type 2 diabetes: What Are They Based on? *Journal of Nutrition and Metabolism*, 2012, 1–6. <https://doi.org/10.1155/2012/847202>

Church, T. S., Blair, S. N., Cocroham, S., Johannsen, N., Johnson, W., Kramer, K., Mikus, C. R., Myers, V., Nauta, M., Rodarte, R. Q., Sparks, L., Thompson, A., & Earnest, C. P. (2010). Effects of Aerobic and Resistance Training on Hemoglobin A_{1c} Levels in Patients With Type 2 Diabetes. *JAMA*, 304(20). <https://doi.org/10.1001/jama.2010.1710>

Clinical applicability and cost-effectiveness of DIABSCORE in screening for type 2 diabetes in primary care- *ClinicalKey*. (n.d.).
<https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S0168822717301031>

Colberg, S. R., Sigal, R. J., Fernhall, B., Regensteiner, J. G., Blissmer, B. J., Rubin, R. R., Chasan-Taber, L., Albright, A. L., & Braun, B. (2010). Exercise and Type 2 Diabetes: The American College of Sports Medicine and the American Diabetes Association: joint position statement executive summary. *Diabetes Care*, 33(12), 2692–2696.
<https://doi.org/10.2337/dc10-1548>

Colberg, S. R., Sigal, R. J., Yardley, J. E., Riddell, M. C., Dunstan, D. W., Dempsey, P. C., Horton, E. S., Castorino, K., & Tate, D. F. (2016). Physical Activity/Exercise and Diabetes: A Position Statement of the American Diabetes Association. *Diabetes Care*, 39(11), 2065–2079. <https://doi.org/10.2337/dc16-1728>

Correction. (2007). *Circulation*, 116(23), e557–e557.
<https://doi.org/10.1161/CIRCULATIONAHA.107.187928>

Creme, P., & Lea, M. R. (2008). *Writing At University: A Guide For Students* (3rd ed). McGraw-Hill Education.
<https://ebookcentral.proquest.com/lib/leicester/detail.action?pq-origsite=primo&docID=345134>

Dagfinn Aune. (2016). Whole grain consumption and risk of cardiovascular disease, cancer, and all cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies. *The BMJ*, 353. <https://doi.org/doi:10.1136/bmj.i2716>

Deborah F Tate. (2012). Replacing caloric beverages with water or diet beverages for weight loss in adults: main results of the Choose Healthy Options Consciously Everyday (CHOICE) randomized clinical trial. *The American Journal of Clinical Nutrition*, 95(3).
<https://doi.org/doi:10.3945/ajcn.111.026278>

Dhingra, R., Sullivan, L., Jacques, P. F., Wang, T. J., Fox, C. S., Meigs, J. B., D'Agostino, R. B., Gaziano, J. M., & Vasan, R. S. (2007). Soft Drink Consumption and Risk of Developing Cardiometabolic Risk Factors and the Metabolic Syndrome in Middle-Aged Adults in the Community. *Circulation*, 116(5), 480–488.
<https://doi.org/10.1161/CIRCULATIONAHA.107.689935>

Differences in the prospective association between individual plasma phospholipid saturated fatty acids and incident type 2 diabetes: the EPIC-InterAct case-cohort study. (n.d.).
http://ac.els-cdn.com/S2213858714701469/1-s2.0-S2213858714701469-main.pdf?_tid=41970bbc-7e93-11e7-a8e5-00000aab0f01&acdnat=1502455838_1e4678aef6e5b79c43c42f6b68f370f1

Dyson, P. A., Kelly, T., Deakin, T., Duncan, A., Frost, G., Harrison, Z., Khatri, D., Kunka, D., McArdle, P., Mellor, D., Oliver, L., & Worth, J. (2011). Diabetes UK evidence-based nutrition guidelines for the prevention and management of diabetes. *Diabetic Medicine*, 28(11), 1282–1288. <https://doi.org/10.1111/j.1464-5491.2011.03371.x>

Effect of a lifestyle intervention on weight change in south Asian individuals in the UK at high risk of type 2 diabetes: a family-cluster randomised controlled trial- *ClinicalKey*. (n.d.).
<https://www.clinicalkey.com/#!/content/journal/1-s2.0-S2213858713702043>

Emily A Hu. (2012). White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review. *The BMJ*, 344. <https://doi.org/doi:10.1136/bmj.e1454>

Fogelholm, M., Larsen, T., Westerterp-Plantenga, M., Macdonald, I., Martinez, J., Boyadjieva, N., Poppitt, S., Schlicht, W., Stratton, G., Sundvall, J., Lam, T., Jalo, E., Christensen, P., Drummen, M., Simpson, E., Navas-Carretero, S., Handjieva-Darlenska, T.,

- Muirhead, R., Silvestre, M., ... Raben, A. (2017). PREVIEW: Prevention of Diabetes through Lifestyle Intervention and Population Studies in Europe and around the World. Design, Methods, and Baseline Participant Description of an Adult Cohort Enrolled into a Three-Year Randomised Clinical Trial. *Nutrients*, 9(6). <https://doi.org/10.3390/nu9060632>
- Georgios S Papaetis. (2014). Incretin-based therapies in prediabetes: Current evidence and future perspectives. *World Journal of Diabetes*, 5(6). <https://doi.org/doi:10.4239/wjd.v5.i6.817>
- Gillies, C. L., Abrams, K. R., Lambert, P. C., Cooper, N. J., Sutton, A. J., Hsu, R. T., & Khunti, K. (2007). Pharmacological and lifestyle interventions to prevent or delay type 2 diabetes in people with impaired glucose tolerance: systematic review and meta-analysis. *BMJ*, 334(7588), 299–299. <https://doi.org/10.1136/bmj.39063.689375.55>
- Gopalan, A., Lorincz, I. S., Wirtalla, C., Marcus, S. C., & Long, J. A. (2015). Awareness of Prediabetes and Engagement in Diabetes Risk-Reducing Behaviors. *American Journal of Preventive Medicine*, 49(4), 512–519. <https://doi.org/10.1016/j.amepre.2015.03.007>
- Gray, L. J., Davies, M. J., Hiles, S., Taub, N. A., Webb, D. R., Srinivasan, B. T., & Khunti, K. (2012). Detection of impaired glucose regulation and/or type 2 diabetes mellitus, using primary care electronic data, in a multiethnic UK community setting. *Diabetologia*, 55(4), 959–966. <https://doi.org/10.1007/s00125-011-2432-x>
- Gray, L. J., Taub, N. A., Khunti, K., Gardiner, E., Hiles, S., Webb, D. R., Srinivasan, B. T., & Davies, M. J. (2010). The Leicester Risk Assessment score for detecting undiagnosed Type 2 diabetes and impaired glucose regulation for use in a multiethnic UK setting. *Diabetic Medicine*, 27(8), 887–895. <https://doi.org/10.1111/j.1464-5491.2010.03037.x>
- Gray, L. J., Yates, T., Troughton, J., Khunti, K., & Davies, M. J. (2016). Engagement, Retention, and Progression to Type 2 Diabetes: A Retrospective Analysis of the Cluster-Randomised 'Let's Prevent Diabetes' Trial. *PLOS Medicine*, 13(7). <https://doi.org/10.1371/journal.pmed.1002078>
- Gregg, E. W., & Shaw, J. E. (2017). Global Health Effects of Overweight and Obesity. *New England Journal of Medicine*. <https://doi.org/10.1056/NEJMe1706095>
- Hamman, R. F., Wing, R. R., Edelstein, S. L., Lachin, J. M., Bray, G. A., Delahanty, L., Hoskin, M., Kriska, A. M., Mayer-Davis, E. J., Pi-Sunyer, X., Regensteiner, J., Venditti, B., & Wylie-Rosett, J. (2006). Effect of Weight Loss With Lifestyle Intervention on Risk of Diabetes. *Diabetes Care*, 29(9), 2102–2107. <https://doi.org/10.2337/dc06-0560>
- Health Effects of Overweight and Obesity in 195 Countries over 25 Years. (2017). *New England Journal of Medicine*. <https://doi.org/10.1056/NEJMoa1614362>
- Helen C Eborall. (2007). Psychological impact of screening for type 2 diabetes: controlled trial and comparative study embedded in the ADDITION (Cambridge) randomised controlled trial. *BMJ: British Medical Journal*, 335(7618). <https://doi.org/doi:10.1136/bmj.39303.723449.55>
- How to prevent type 2 diabetes in women with previous gestational diabetes? A systematic review of behavioural interventions- ClinicalKey. (n.d.). <https://www.clinicalkey.com/#!/content/playContent/1-s2.0-S1751991817300785?returnurl>

=null&referrer=null

Joiner, K. L., Nam, S., & Whittemore, R. (2017). Lifestyle interventions based on the diabetes prevention program delivered via eHealth: A systematic review and meta-analysis. *Preventive Medicine*, 100, 194–207.
<https://doi.org/10.1016/j.ypmed.2017.04.033>

Koivusalo, S. B., Rönö, K., Klemetti, M. M., Roine, R. P., Lindström, J., Erkkola, M., Kaaja, R. J., Pöyhönen-Alho, M., Tiitinen, A., Huvinen, E., Andersson, S., Laivuori, H., Valkama, A., Meinilä, J., Kautiainen, H., Eriksson, J. G., & Stach-Lempinen, B. (2016). Gestational Diabetes Mellitus Can Be Prevented by Lifestyle Intervention: The Finnish Gestational Diabetes Prevention Study (RADIEL). *Diabetes Care*, 39(1), 24–30.
<https://doi.org/10.2337/dc15-0511>

Laaksonen, D. E., Lindstrom, J., Lakka, T. A., Eriksson, J. G., Niskanen, L., Wikstrom, K., Aunola, S., Keinänen-Kiukaanniemi, S., Laakso, M., Valle, T. T., Ilanne-Parikka, P., Louheranta, A., Hamalainen, H., Rastas, M., Salminen, V., Cepaitis, Z., Hakumaki, M., Kaikkonen, H., Harkonen, P., ... Uusitupa, M. (2005). Physical Activity in the Prevention of Type 2 Diabetes: The Finnish Diabetes Prevention Study. *Diabetes*, 54(1), 158–165.
<https://doi.org/10.2337/diabetes.54.1.158>

LaMonte, M. J. (2005). Physical activity and diabetes prevention. *Journal of Applied Physiology*, 99(3), 1205–1213. <https://doi.org/10.1152/jappphysiol.00193.2005>

Less Sitting, More Physical Activity, or Higher Fitness?- ClinicalKey. (n.d.).
<https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/playContent/1-s2.0-S0025619615006308?returnurl=null&referrer=null>

Ley, S. H., Hamdy, O., Mohan, V., & Hu, F. B. (2014). Prevention and management of type 2 diabetes: dietary components and nutritional strategies. *The Lancet*, 383(9933), 1999–2007. [https://doi.org/10.1016/S0140-6736\(14\)60613-9](https://doi.org/10.1016/S0140-6736(14)60613-9)

Li, R., Zhang, P., Barker, L. E., Chowdhury, F. M., & Zhang, X. (2010). Cost-Effectiveness of Interventions to Prevent and Control Diabetes Mellitus: A Systematic Review. *Diabetes Care*, 33(8), 1872–1894. <https://doi.org/10.2337/dc10-0843>

Linda Penn. (2013a). Importance of Weight Loss Maintenance and Risk Prediction in the Prevention of Type 2 Diabetes: Analysis of European Diabetes Prevention Study RCT. *PLoS ONE*, 8(2). <https://doi.org/doi:10.1371/journal.pone.0057143>

Linda Penn. (2013b). Importance of Weight Loss Maintenance and Risk Prediction in the Prevention of Type 2 Diabetes: Analysis of European Diabetes Prevention Study RCT. *PLoS ONE*, 8(2). <https://doi.org/doi:10.1371/journal.pone.0057143>

Lindstrom, J., Louheranta, A., Mannelin, M., Rastas, M., Salminen, V., Eriksson, J., Uusitupa, M., & Tuomilehto, J. (2003). The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care*, 26(12), 3230–3236. <https://doi.org/10.2337/diacare.26.12.3230>

Long, G. H., Johansson, I., Rolandsson, O., Wennberg, P., Fhärm, E., Weinehall, L., Griffin, S. J., Simmons, R. K., & Norberg, M. (2015). Healthy behaviours and 10-year incidence of

diabetes: A population cohort study. *Preventive Medicine*, 71, 121–127.
<https://doi.org/10.1016/j.ypmed.2014.12.013>

Luchsinger, J. A., Ma, Y., Christophi, C. A., Florez, H., Golden, S. H., Hazuda, H., Crandall, J., Venditti, E., Watson, K., Jeffries, S., Manly, J. J., & Pi-Sunyer, F. X. (2017). Metformin, Lifestyle Intervention, and Cognition in the Diabetes Prevention Program Outcomes Study. *Diabetes Care*, 40(7), 958–965. <https://doi.org/10.2337/dc16-2376>

Malik, V. S., Pan, A., Willett, W. C., & Hu, F. B. (2013). Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis. *American Journal of Clinical Nutrition*, 98(4), 1084–1102. <https://doi.org/10.3945/ajcn.113.058362>

Malik, V. S., Popkin, B. M., Bray, G. A., Despres, J.-P., Willett, W. C., & Hu, F. B. (2010). Sugar-Sweetened Beverages and Risk of Metabolic Syndrome and Type 2 Diabetes: A meta-analysis. *Diabetes Care*, 33(11), 2477–2483. <https://doi.org/10.2337/dc10-1079>

Mann, J., Morenga, L. T., McLean, R., Swinburn, B., Mhurchu, C. N., Jackson, R., Kennedy, J., & Beaglehole, R. (2016). Dietary guidelines on trial: the charges are not evidence based. *The Lancet*, 388(10047), 851–853. [https://doi.org/10.1016/S0140-6736\(16\)31278-8](https://doi.org/10.1016/S0140-6736(16)31278-8)

Matti Uusitupa. (2009). Ten-Year Mortality and Cardiovascular Morbidity in the Finnish Diabetes Prevention Study—Secondary Analysis of the Randomized Trial. *PLoS ONE*, 4(5). <https://doi.org/doi:10.1371/journal.pone.0005656>

Nathan, D. M., Davidson, M. B., DeFronzo, R. A., Heine, R. J., Henry, R. R., Pratley, R., & Zinman, B. (2007). Impaired Fasting Glucose and Impaired Glucose Tolerance: Implications for care. *Diabetes Care*, 30(3), 753–759. <https://doi.org/10.2337/dc07-9920>

NICE - Type 2 diabetes: prevention in people at high risk | Guidance and guidelines. (n.d.). NICE. <https://www.nice.org.uk/guidance/ph38>

Nutrition Recommendations and Interventions for Diabetes: A position statement of the American Diabetes Association. (2008). *Diabetes Care*, 31(Supplement 1), S61–S78. <https://doi.org/10.2337/dc08-S061>

Obesity: identification, assessment and management | Guidance and guidelines | NICE. (n.d.). <https://www.nice.org.uk/guidance/cg189>

Orozco, L. J., Buchleitner, A. M., Gimenez-Perez, G., Roqué i Figuls, M., Richter, B., & Mauricio, D. (1996). Exercise or exercise and diet for preventing type 2 diabetes mellitus. In *Cochrane Database of Systematic Reviews*. John Wiley & Sons, Ltd. <https://doi.org/10.1002/14651858.CD003054.pub3>

Pan, A., Sun, Q., Bernstein, A. M., Schulze, M. B., Manson, J. E., Willett, W. C., & Hu, F. B. (2011). Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis. *American Journal of Clinical Nutrition*, 94(4), 1088–1096. <https://doi.org/10.3945/ajcn.111.018978>

Pan, X.-R., Li, G.-W., Hu, Y.-H., Wang, J.-X., Yang, W.-Y., An, Z.-X., Hu, Z.-X., Xiao, J.-Z., Cao, H.-B., Liu, P.-A., Jiang, X.-G., Jiang, Y.-Y., Wang, J.-P., Zheng, H., Zhang, H., Bennett, P. H., & Howard, B. V. (1997). Effects of Diet and Exercise in Preventing NIDDM in People With

Impaired Glucose Tolerance: The Da Qing IGT and Diabetes Study. *Diabetes Care*, 20(4), 537–544. <https://doi.org/10.2337/diacare.20.4.537>

Public Health England - Diabetes prevention programmes: evidence review. (2015). Public Health England.
<https://www.gov.uk/government/publications/diabetes-prevention-programmes-evidence-review>

Rajul Parikh, Annie Mathai, Shefali Parikh, G. Chandra Sekhar, Ravi Thomas. (n.d.). Understanding and using sensitivity, specificity and predictive values. *Indian Journal of Ophthalmology*.
http://go.galegroup.com.ezproxy4.lib.le.ac.uk/ps/retrieve.do?tabID=T002&resultListType=RESULT_LIST&searchResultsType=SingleTab&searchType=AdvancedSearchForm¤tPosition=3&docId=GALE%7CA173264871&docType=Article&sort=Relevance&contentSegment=&prodId=EAIM&contentSet=GALE%7CA173264871&searchId=R1&userGroupName=leicester&inPS=true

Ramachandran, A., Snehalatha, C., Mary, S., Mukesh, B., Bhaskar, A. D., & Vijay, V. (2006). The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia*, 49(2), 289–297.
<https://doi.org/10.1007/s00125-005-0097-z>

Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin. (2002a). *New England Journal of Medicine*, 346(6), 393–403.
<https://doi.org/10.1056/NEJMoa012512>

Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin. (2002b). *New England Journal of Medicine*, 346(6), 393–403.
<https://doi.org/10.1056/NEJMoa012512>

Schulze, M. B. (2004). Sugar-Sweetened Beverages, Weight Gain, and Incidence of Type 2 Diabetes in Young and Middle-Aged Women. *JAMA*, 292(8).
<https://doi.org/10.1001/jama.292.8.927>

Schwingshackl, L., Hoffmann, G., Lampousi, A.-M., Knüppel, S., Iqbal, K., Schwedhelm, C., Bechthold, A., Schlesinger, S., & Boeing, H. (2017). Food groups and risk of type 2 diabetes mellitus: a systematic review and meta-analysis of prospective studies. *European Journal of Epidemiology*, 32(5), 363–375. <https://doi.org/10.1007/s10654-017-0246-y>

Screening for type 2 diabetes: literature review and economic modelling. (2007). *Clinical Governance: An International Journal*, 12(4).
<https://doi.org/10.1108/cgij.2007.24812dae.003>

Sedentary Behaviors and Subsequent Health Outcomes in Adults- ClinicalKey. (n.d.-a).
<https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/journal/1-s2.0-S0749379711003126>

Sedentary Behaviors and Subsequent Health Outcomes in Adults- ClinicalKey. (n.d.-b).
<https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/journal/1-s2.0-S0749379711003126>

Stefan, N., Staiger, H., Wagner, R., Machann, J., Schick, F., H?ring, H.-U., & Fritsche, A. (2015). A high-risk phenotype associates with reduced improvement in glycaemia during a lifestyle intervention in prediabetes. *Diabetologia*, 58(12), 2877–2884. <https://doi.org/10.1007/s00125-015-3760-z>

Steven N Blair. (2009). Physical inactivity: the biggest public health problem of the 21st century. *British Journal of Sports Medicine*, 43, 1–2. <http://bjsm.bmj.com/content/43/1/1.full>

Sweeting, A. N., Ross, G. P., Hyett, J., Molyneaux, L., Constantino, M., Harding, A. J., & Wong, J. (2016). Gestational Diabetes Mellitus in Early Pregnancy: Evidence for Poor Pregnancy Outcomes Despite Treatment. *Diabetes Care*, 39(1), 75–81. <https://doi.org/10.2337/dc15-0433>

The 10-Year Cost-Effectiveness of Lifestyle Intervention or Metformin for Diabetes Prevention: An intent-to-treat analysis of the DPP/DPPOS. (2012). *Diabetes Care*, 35(4), 723–730. <https://doi.org/10.2337/dc11-1468>

Thomas Yates. (2012). Walking away from type 2 diabetes: trial protocol of a cluster randomised controlled trial evaluating a structured education programme in those at high risk of developing type 2 diabetes. *BMC Family Practice*, 13. <https://doi.org/doi:10.1186/1471-2296-13-46>

Tseng, E., Yeh, H.-C., & Maruthur, N. M. (2017). Metformin Use in Prediabetes Among U.S. Adults, 2005?2012. *Diabetes Care*, 40(7), 887–893. <https://doi.org/10.2337/dc16-1509>

Tuomilehto, J., Lindstrom, J., Eriksson, J. G., Valle, T. T., Hamalainen, H., Ilanne-Parikka, P., Keinanen-Kiukaanniemi, S., Laakso, M., Louheranta, A., Rastas, M., Salminen, V., Aunola, S., Cepaitis, Z., Moltchanov, V., Hakumaki, M., Mannelin, M., Martikkala, V., Sundvall, J., & Uusitupa, M. (2001). Prevention of Type 2 Diabetes Mellitus by Changes in Lifestyle among Subjects with Impaired Glucose Tolerance. *New England Journal of Medicine*, 344(18), 1343–1350. <https://doi.org/10.1056/NEJM200105033441801>

Type 2 diabetes: prevention in people at high risk | Guidance and guidelines | NICE. (n.d.). <https://www.nice.org.uk/guidance/ph38/documents/preventing-type-2-diabetes-risk-identification-and-interventions-for-individuals-at-high-risk-draft-guidance-consultation>

Vita, P., Cardona-Morrell, M., Bauman, A., Singh, M. F., Moore, M., Pennock, R., Snow, J., Williams, M., Jackson, L., Milat, A., & Colagiuri, S. (2016). Type 2 diabetes prevention in the community: 12-Month outcomes from the Sydney Diabetes Prevention Program. *Diabetes Research and Clinical Practice*, 112, 13–19. <https://doi.org/10.1016/j.diabres.2015.11.010>
Webb, D. R., Gray, L. J., Khunti, K., Srinivasan, B., Taub, N., Campbell, S., Barnett, J., Farooqi, A., Echouffo-Tcheugui, J. B., Griffin, S. J., Wareham, N. J., & Davies, M. J. (2011). Screening for diabetes using an oral glucose tolerance test within a western multi-ethnic population identifies modifiable cardiovascular risk: the ADDITION-Leicester study. *Diabetologia*, 54(9), 2237–2246. <https://doi.org/10.1007/s00125-011-2189-2>

Wu, Y., Zhang, D., Jiang, X., & Jiang, W. (2015). Fruit and vegetable consumption and risk of type 2 diabetes mellitus: A dose-response meta-analysis of prospective cohort studies. *Nutrition, Metabolism and Cardiovascular Diseases*, 25(2), 140–147. <https://doi.org/10.1016/j.numecd.2014.10.004>

Yates, T., Wilmot, E. G., Khunti, K., Biddle, S., Gorely, T., & Davies, M. J. (2011). Stand up for your health: Is it time to rethink the physical activity paradigm? *Diabetes Research and Clinical Practice*, 93(2), 292–294. <https://doi.org/10.1016/j.diabres.2011.03.023>