

MD7513 - Prevention, Screening & Early Detection in Diabetes

[View Online](#)

Prevention, Screening & Early Detection in Diabetes

[1]

'Public Health England - Diabetes prevention programmes: evidence review'. Public Health England, 2015. [Online]. Available: <https://www.gov.uk/government/publications/diabetes-prevention-programmes-evidence-review>

[2]

'American Diabetes Association - Standards of Medical Care in Diabetes - 2017 Summary of Revisions', Diabetes Care, vol. 40, no. Supplement 1, pp. S4-S5, Jan. 2017, doi: 10.2337/dc17-S003.

[3]

'NICE - Type 2 diabetes: prevention in people at high risk | Guidance and guidelines'. NICE. [Online]. Available: <https://www.nice.org.uk/guidance/ph38>

[4]

'Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin', New England Journal of Medicine, vol. 346, no. 6, pp. 393-403, Feb. 2002, doi: 10.1056/NEJMoa012512.

[5]

C. L. Gillies et al., 'Pharmacological and lifestyle interventions to prevent or delay type 2 diabetes in people with impaired glucose tolerance: systematic review and meta-analysis', BMJ, vol. 334, no. 7588, pp. 299-299, Jan. 2007, doi: 10.1136/bmj.39063.689375.55.

[6]

J. Lindstrom et al., 'The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity', *Diabetes Care*, vol. 26, no. 12, pp. 3230–3236, Dec. 2003, doi: 10.2337/diacare.26.12.3230.

[7]

A. Ramachandran, C. Snehalatha, S. Mary, B. Mukesh, A. D. Bhaskar, and V. Vijay, 'The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1)', *Diabetologia*, vol. 49, no. 2, pp. 289–297, Feb. 2006, doi: 10.1007/s00125-005-0097-z.

[8]

J. Tuomilehto et al., 'Prevention of Type 2 Diabetes Mellitus by Changes in Lifestyle among Subjects with Impaired Glucose Tolerance', *New England Journal of Medicine*, vol. 344, no. 18, pp. 1343–1350, May 2001, doi: 10.1056/NEJM200105033441801.

[9]

Matti Uusitupa, 'Ten-Year Mortality and Cardiovascular Morbidity in the Finnish Diabetes Prevention Study—Secondary Analysis of the Randomized Trial', *PLoS ONE*, vol. 4, no. 5, 2009, doi: 10.1371/journal.pone.0005656.

[10]

J. Brown et al., 'Lifestyle interventions for the treatment of women with gestational diabetes', in *Cochrane Database of Systematic Reviews*, Chichester, UK: John Wiley & Sons, Ltd, 1996. doi: 10.1002/14651858.CD011970.pub2.

[11]

L. M. S. Carlsson et al., 'Bariatric Surgery and Prevention of Type 2 Diabetes in Swedish Obese Subjects', *New England Journal of Medicine*, vol. 367, no. 8, pp. 695–704, Aug. 2012, doi: 10.1056/NEJMoa1112082.

[12]

E. Tseng, H.-C. Yeh, and N. M. Maruthur, 'Metformin Use in Prediabetes Among U.S. Adults, 2005-2012', *Diabetes Care*, vol. 40, no. 7, pp. 887-893, Jul. 2017, doi: 10.2337/dc16-1509.

[13]

J. A. Luchsinger et al., 'Metformin, Lifestyle Intervention, and Cognition in the Diabetes Prevention Program Outcomes Study', *Diabetes Care*, vol. 40, no. 7, pp. 958-965, Jul. 2017, doi: 10.2337/dc16-2376.

[14]

P. R. Breeze et al., 'The impact of Type 2 diabetes prevention programmes based on risk-identification and lifestyle intervention intensity strategies: a cost-effectiveness analysis', *Diabetic Medicine*, vol. 34, no. 5, pp. 632-640, May 2017, doi: 10.1111/dme.13314.

[15]

M. Fogelholm et al., 'PREVIEW: Prevention of Diabetes through Lifestyle Intervention and Population Studies in Europe and around the World. Design, Methods, and Baseline Participant Description of an Adult Cohort Enrolled into a Three-Year Randomised Clinical Trial', *Nutrients*, vol. 9, no. 6, Jun. 2017, doi: 10.3390/nu9060632.

[16]

'Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin', *New England Journal of Medicine*, vol. 346, no. 6, pp. 393-403, Feb. 2002, doi: 10.1056/NEJMoa012512.

[17]

L. J. Gray, T. Yates, J. Troughton, K. Khunti, and M. J. Davies, 'Engagement, Retention, and Progression to Type 2 Diabetes: A Retrospective Analysis of the Cluster-Randomised "Let's Prevent Diabetes" Trial', *PLOS Medicine*, vol. 13, no. 7, Jul. 2016, doi: 10.1371/journal.pmed.1002078.

[18]

'A community based primary prevention programme for type 2 diabetes integrating identification and lifestyle intervention for prevention: the Let's Prevent Diabetes cluster randomised controlled trial- ClinicalKey'. [Online]. Available: <https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S0091743515003850>

[19]

E. Barry, S. Roberts, J. Oke, S. Vijayaraghavan, R. Normansell, and T. Greenhalgh, 'Efficacy and effectiveness of screen and treat policies in prevention of type 2 diabetes: systematic review and meta-analysis of screening tests and interventions', *BMJ*, Jan. 2017, doi: 10.1136/bmj.i6538.

[20]

D. R. Webb et al., 'Screening for diabetes using an oral glucose tolerance test within a western multi-ethnic population identifies modifiable cardiovascular risk: the ADDITION-Leicester study', *Diabetologia*, vol. 54, no. 9, pp. 2237–2246, Sep. 2011, doi: 10.1007/s00125-011-2189-2.

[21]

'Screening for type 2 diabetes: literature review and economic modelling', *Clinical Governance: An International Journal*, vol. 12, no. 4, Oct. 2007, doi: 10.1108/cgij.2007.24812dae.003.

[22]

L. J. Gray et al., 'The Leicester Risk Assessment score for detecting undiagnosed Type 2 diabetes and impaired glucose regulation for use in a multiethnic UK setting', *Diabetic Medicine*, vol. 27, no. 8, pp. 887–895, May 2010, doi: 10.1111/j.1464-5491.2010.03037.x.

[23]

L. J. Gray et al., 'Detection of impaired glucose regulation and/or type 2 diabetes mellitus, using primary care electronic data, in a multiethnic UK community setting', *Diabetologia*, vol. 55, no. 4, pp. 959–966, Apr. 2012, doi: 10.1007/s00125-011-2432-x.

[24]

Rajul Parikh, Annie Mathai, Shefali Parikh, G. Chandra Sekhar, Ravi Thomas, 'Understanding and using sensitivity, specificity and predictive values', Indian Journal of Ophthalmology, [Online]. Available:
http://go.galegroup.com.ezproxy4.lib.le.ac.uk/ps/retrieve.do?tabID=T002&resultListType=RESULT_LIST&searchResultsType=SingleTab&searchType=AdvancedSearchForm¤tPosition=3&docId=GALE%7CA173264871&docType=Article&sort=Relevance&contentSegment=&prodId=EAIM&contentSet=GALE%7CA173264871&searchId=R1&userGroupName=leicester&inPS=true

[25]

Helen C Eborall, 'Psychological impact of screening for type 2 diabetes: controlled trial and comparative study embedded in the ADDITION (Cambridge) randomised controlled trial', BMJ : British Medical Journal, vol. 335, no. 7618, 2007, doi: doi: 10.1136/bmj.39303.723449.55.

[26]

R. Li, P. Zhang, L. E. Barker, F. M. Chowdhury, and X. Zhang, 'Cost-Effectiveness of Interventions to Prevent and Control Diabetes Mellitus: A Systematic Review', Diabetes Care, vol. 33, no. 8, pp. 1872–1894, Aug. 2010, doi: 10.2337/dc10-0843.

[27]

'Clinical applicability and cost-effectiveness of DIABSCORE in screening for type 2 diabetes in primary care- ClinicalKey'. [Online]. Available:
<https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S0168822717301031>

[28]

R. F. Hamman et al., 'Effect of Weight Loss With Lifestyle Intervention on Risk of Diabetes', Diabetes Care, vol. 29, no. 9, pp. 2102–2107, Sep. 2006, doi: 10.2337/dc06-0560.

[29]

K. L. Joiner, S. Nam, and R. Whittemore, 'Lifestyle interventions based on the diabetes

prevention program delivered via eHealth: A systematic review and meta-analysis', Preventive Medicine, vol. 100, pp. 194–207, Jul. 2017, doi: 10.1016/j.ypmed.2017.04.033.

[30]

Georgios S Papaetis, 'Incretin-based therapies in prediabetes: Current evidence and future perspectives', World Journal of Diabetes, vol. 5, no. 6, 2014, doi: doi: 10.4239/wjd.v5.i6.817.

[31]

'Obesity: identification, assessment and management | Guidance and guidelines | NICE', [Online]. Available: <https://www.nice.org.uk/guidance/cg189>

[32]

'Health Effects of Overweight and Obesity in 195 Countries over 25 Years', New England Journal of Medicine, Jun. 2017, doi: 10.1056/NEJMoa1614362.

[33]

E. W. Gregg and J. E. Shaw, 'Global Health Effects of Overweight and Obesity', New England Journal of Medicine, Jun. 2017, doi: 10.1056/NEJMe1706095.

[34]

'A Systems Science Approach to the Obesity Epidemic and Prevention of Diabetes.' 20 AD. [Online]. Available: <https://www.youtube.com/watch?v=HrrD-Z0B7dY&feature=youtu.be&t=55s%20A%20Syst%20Science%20Approach%20to%20the%20Obesity%20Epidemic%20and%20Prevention%20of%20Diabetes>.

[35]

L. M. S. Carlsson et al., 'Bariatric Surgery and Prevention of Type 2 Diabetes in Swedish Obese Subjects', New England Journal of Medicine, vol. 367, no. 8, pp. 695–704, Aug. 2012, doi: 10.1056/NEJMoa1112082.

[36]

Linda Penn, 'Importance of Weight Loss Maintenance and Risk Prediction in the Prevention of Type 2 Diabetes: Analysis of European Diabetes Prevention Study RCT', PLoS ONE, vol. 8, no. 2, 2013, doi: doi: 10.1371/journal.pone.0057143.

[37]

E. J. Aguiar, P. J. Morgan, C. E. Collins, R. C. Plotnikoff, M. D. Young, and R. Callister, 'Efficacy of the Type 2 Diabetes Prevention Using LifeStyle Education Program RCT', American Journal of Preventive Medicine, vol. 50, no. 3, pp. 353–364, Mar. 2016, doi: 10.1016/j.amepre.2015.08.020.

[38]

'Effect of a lifestyle intervention on weight change in south Asian individuals in the UK at high risk of type 2 diabetes: a family-cluster randomised controlled trial- ClinicalKey'. [Online]. Available: <https://www.clinicalkey.com/#!/content/journal/1-s2.0-S2213858713702043>

[39]

Linda Penn, 'Importance of Weight Loss Maintenance and Risk Prediction in the Prevention of Type 2 Diabetes: Analysis of European Diabetes Prevention Study RCT', PLoS ONE, vol. 8, no. 2, 2013, doi: doi: 10.1371/journal.pone.0057143.

[40]

G. H. Long et al., 'Healthy behaviours and 10-year incidence of diabetes: A population cohort study', Preventive Medicine, vol. 71, pp. 121–127, Feb. 2015, doi: 10.1016/j.ypmed.2014.12.013.

[41]

'Nutrition Recommendations and Interventions for Diabetes: A position statement of the American Diabetes Association', Diabetes Care, vol. 31, no. Supplement 1, pp. S61–S78, Jan. 2008, doi: 10.2337/dc08-S061.

[42]

P. Carter, K. Khunti, and M. J. Davies, 'Dietary Recommendations for the Prevention of Type 2 diabetes: What Are They Based on?', *Journal of Nutrition and Metabolism*, vol. 2012, pp. 1–6, 2012, doi: 10.1155/2012/847202.

[43]

X.-R. Pan et al., 'Effects of Diet and Exercise in Preventing NIDDM in People With Impaired Glucose Tolerance: The Da Qing IGT and Diabetes Study', *Diabetes Care*, vol. 20, no. 4, pp. 537–544, Apr. 1997, doi: 10.2337/diacare.20.4.537.

[44]

'How to prevent type 2 diabetes in women with previous gestational diabetes? A systematic review of behavioural interventions- ClinicalKey'. [Online]. Available: <https://www.clinicalkey.com/#!/content/playContent/1-s2.0-S1751991817300785?returnurl=null&referrer=null>

[45]

V. S. Malik, B. M. Popkin, G. A. Bray, J.-P. Despres, W. C. Willett, and F. B. Hu, 'Sugar-Sweetened Beverages and Risk of Metabolic Syndrome and Type 2 Diabetes: A meta-analysis', *Diabetes Care*, vol. 33, no. 11, pp. 2477–2483, Nov. 2010, doi: 10.2337/dc10-1079.

[46]

M. B. Schulze, 'Sugar-Sweetened Beverages, Weight Gain, and Incidence of Type 2 Diabetes in Young and Middle-Aged Women', *JAMA*, vol. 292, no. 8, Aug. 2004, doi: 10.1001/jama.292.8.927.

[47]

'Correction', *Circulation*, vol. 116, no. 23, pp. e557–e557, Dec. 2007, doi: 10.1161/CIRCULATIONAHA.107.187928.

[48]

V. S. Malik, A. Pan, W. C. Willett, and F. B. Hu, 'Sugar-sweetened beverages and weight gain in children and adults: a systematic review and meta-analysis', *American Journal of Clinical Nutrition*, vol. 98, no. 4, pp. 1084–1102, Oct. 2013, doi: 10.3945/ajcn.113.058362.

[49]

Deborah F Tate, 'Replacing caloric beverages with water or diet beverages for weight loss in adults: main results of the Choose Healthy Options Consciously Everyday (CHOICE) randomized clinical trial', *The American Journal of Clinical Nutrition*, vol. 95, no. 3, 2012, doi: 10.3945/ajcn.111.026278.

[50]

A. Pan et al., 'Red meat consumption and risk of type 2 diabetes: 3 cohorts of US adults and an updated meta-analysis', *American Journal of Clinical Nutrition*, vol. 94, no. 4, pp. 1088–1096, Oct. 2011, doi: 10.3945/ajcn.111.018978.

[51]

S. H. Ley, O. Hamdy, V. Mohan, and F. B. Hu, 'Prevention and management of type 2 diabetes: dietary components and nutritional strategies', *The Lancet*, vol. 383, no. 9933, pp. 1999–2007, Jun. 2014, doi: 10.1016/S0140-6736(14)60613-9.

[52]

'Type 2 diabetes: prevention in people at high risk | Guidance and guidelines | NICE', [Online]. Available: <https://www.nice.org.uk/guidance/ph38/documents/preventing-type-2-diabetes-risk-identification-and-interventions-for-individuals-at-high-risk-draft-guidance-consultation>

[53]

Emily A Hu, 'White rice consumption and risk of type 2 diabetes: meta-analysis and systematic review', *The BMJ*, vol. 344, 2012, doi: 10.1136/bmj.e1454.

[54]

Dagfinn Aune, 'Whole grain consumption and risk of cardiovascular disease, cancer, and all

cause and cause specific mortality: systematic review and dose-response meta-analysis of prospective studies', *The BMJ*, vol. 353, 2016, doi: doi: 10.1136/bmj.i2716.

[55]

P. A. Dyson et al., 'Diabetes UK evidence-based nutrition guidelines for the prevention and management of diabetes', *Diabetic Medicine*, vol. 28, no. 11, pp. 1282–1288, Nov. 2011, doi: 10.1111/j.1464-5491.2011.03371.x.

[56]

L. Schwingshackl et al., 'Food groups and risk of type 2 diabetes mellitus: a systematic review and meta-analysis of prospective studies', *European Journal of Epidemiology*, vol. 32, no. 5, pp. 363–375, May 2017, doi: 10.1007/s10654-017-0246-y.

[57]

J. Mann et al., 'Dietary guidelines on trial: the charges are not evidence based', *The Lancet*, vol. 388, no. 10047, pp. 851–853, Aug. 2016, doi: 10.1016/S0140-6736(16)31278-8.

[58]

R. Dhingra et al., 'Soft Drink Consumption and Risk of Developing Cardiometabolic Risk Factors and the Metabolic Syndrome in Middle-Aged Adults in the Community', *Circulation*, vol. 116, no. 5, pp. 480–488, Jul. 2007, doi: 10.1161/CIRCULATIONAHA.107.689935.

[59]

Y. Wu, D. Zhang, X. Jiang, and W. Jiang, 'Fruit and vegetable consumption and risk of type 2 diabetes mellitus: A dose-response meta-analysis of prospective cohort studies', *Nutrition, Metabolism and Cardiovascular Diseases*, vol. 25, no. 2, pp. 140–147, Feb. 2015, doi: 10.1016/j.numecd.2014.10.004.

[60]

'Differences in the prospective association between individual plasma phospholipid saturated fatty acids and incident type 2 diabetes: the EPIC-InterAct case-cohort study'. [Online]. Available: http://ac.els-cdn.com/S2213858714701469/1-s2.0-S2213858714701469-main.pdf?_tid=41

970bbc-7e93-11e7-a8e5-00000aab0f01&acdnat=1502455838_1e4678aef6e5b79c43c42f6b68f370f1

[61]

D. Aune, T. Norat, P. Romundstad, and L. J. Vatten, 'Whole grain and refined grain consumption and the risk of type 2 diabetes: a systematic review and dose-response meta-analysis of cohort studies', *European Journal of Epidemiology*, vol. 28, no. 11, pp. 845–858, Nov. 2013, doi: 10.1007/s10654-013-9852-5.

[62]

S. R. Colberg et al., 'Exercise and Type 2 Diabetes: The American College of Sports Medicine and the American Diabetes Association: joint position statement executive summary', *Diabetes Care*, vol. 33, no. 12, pp. 2692–2696, Dec. 2010, doi: 10.2337/dc10-1548.

[63]

T. Yates, E. G. Wilmot, K. Khunti, S. Biddle, T. Gorely, and M. J. Davies, 'Stand up for your health: Is it time to rethink the physical activity paradigm?', *Diabetes Research and Clinical Practice*, vol. 93, no. 2, pp. 292–294, Aug. 2011, doi: 10.1016/j.diabres.2011.03.023.

[64]

Steven N Blair, 'Physical inactivity: the biggest public health problem of the 21st century', *British Journal of Sports Medicine*, vol. 43, pp. 1–2, Jan. 2009, [Online]. Available: <http://bjsm.bmj.com/content/43/1/1.full>

[65]

D. E. Laaksonen et al., 'Physical Activity in the Prevention of Type 2 Diabetes: The Finnish Diabetes Prevention Study', *Diabetes*, vol. 54, no. 1, pp. 158–165, Jan. 2005, doi: 10.2337/diabetes.54.1.158.

[66]

Thomas Yates, 'Walking away from type 2 diabetes: trial protocol of a cluster randomised controlled trial evaluating a structured education programme in those at high risk of

developing type 2 diabetes', BMC Family Practice, vol. 13, 2012, doi: doi: 10.1186/1471-2296-13-46.

[67]

S. R. Colberg et al., 'Physical Activity/Exercise and Diabetes: A Position Statement of the American Diabetes Association', Diabetes Care, vol. 39, no. 11, pp. 2065–2079, Nov. 2016, doi: 10.2337/dc16-1728.

[68]

N. Stefan et al., 'A high-risk phenotype associates with reduced improvement in glycaemia during a lifestyle intervention in prediabetes', Diabetologia, vol. 58, no. 12, pp. 2877–2884, Dec. 2015, doi: 10.1007/s00125-015-3760-z.

[69]

'Less Sitting, More Physical Activity, or Higher Fitness?- ClinicalKey'. [Online]. Available: <https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/playContent/1-s2.0-S0025619615006308?returnurl=null&referrer=null>

[70]

'Sedentary Behaviors and Subsequent Health Outcomes in Adults- ClinicalKey'. [Online]. Available: <https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/journal/1-s2.0-S0749379711003126>

[71]

M. J. LaMonte, 'Physical activity and diabetes prevention', Journal of Applied Physiology, vol. 99, no. 3, pp. 1205–1213, Sep. 2005, doi: 10.1152/japplphysiol.00193.2005.

[72]

T. S. Church et al., 'Effects of Aerobic and Resistance Training on Hemoglobin A Levels in Patients With Type 2 Diabetes', JAMA, vol. 304, no. 20, Nov. 2010, doi: 10.1001/jama.2010.1710.

[73]

L. J. Orozco, A. M. Buchleitner, G. Gimenez-Perez, M. Roqué i Figuls, B. Richter, and D. Mauricio, 'Exercise or exercise and diet for preventing type 2 diabetes mellitus', in Cochrane Database of Systematic Reviews, Chichester, UK: John Wiley & Sons, Ltd, 1996. doi: 10.1002/14651858.CD003054.pub3.

[74]

'Sedentary Behaviors and Subsequent Health Outcomes in Adults- ClinicalKey'. [Online]. Available:
<https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/journal/1-s2.0-S0749379711003126>

[75]

A. Gopalan, I. S. Lorincz, C. Wirtalla, S. C. Marcus, and J. A. Long, 'Awareness of Prediabetes and Engagement in Diabetes Risk-Reducing Behaviors', American Journal of Preventive Medicine, vol. 49, no. 4, pp. 512–519, Oct. 2015, doi: 10.1016/j.amepre.2015.03.007.

[76]

P. Absetz et al., 'Type 2 Diabetes Prevention in the Real World: Three-year results of the GOAL Lifestyle Implementation Trial', Diabetes Care, vol. 32, no. 8, pp. 1418–1420, Aug. 2009, doi: 10.2337/dc09-0039.

[77]

R. T. Ackermann, E. A. Finch, E. Brizendine, H. Zhou, and D. G. Marrero, 'Translating the Diabetes Prevention Program into the Community', American Journal of Preventive Medicine, vol. 35, no. 4, pp. 357–363, Oct. 2008, doi: 10.1016/j.amepre.2008.06.035.

[78]

A. N. Sweeting et al., 'Gestational Diabetes Mellitus in Early Pregnancy: Evidence for Poor Pregnancy Outcomes Despite Treatment', Diabetes Care, vol. 39, no. 1, pp. 75–81, Jan. 2016, doi: 10.2337/dc15-0433.

[79]

S. B. Koivusalo et al., 'Gestational Diabetes Mellitus Can Be Prevented by Lifestyle Intervention: The Finnish Gestational Diabetes Prevention Study (RADIEL)', *Diabetes Care*, vol. 39, no. 1, pp. 24–30, Jan. 2016, doi: 10.2337/dc15-0511.

[80]

D. M. Nathan et al., 'Impaired Fasting Glucose and Impaired Glucose Tolerance: Implications for care', *Diabetes Care*, vol. 30, no. 3, pp. 753–759, Mar. 2007, doi: 10.2337/dc07-9920.

[81]

P. Vita et al., 'Type 2 diabetes prevention in the community: 12-Month outcomes from the Sydney Diabetes Prevention Program', *Diabetes Research and Clinical Practice*, vol. 112, pp. 13–19, Feb. 2016, doi: 10.1016/j.diabres.2015.11.010.

[82]

P. R. Breeze et al., 'Cost-effectiveness of population-based, community, workplace and individual policies for diabetes prevention in the UK', *Diabetic Medicine*, vol. 34, no. 8, pp. 1136–1144, Aug. 2017, doi: 10.1111/dme.13349.

[83]

'The 10-Year Cost-Effectiveness of Lifestyle Intervention or Metformin for Diabetes Prevention: An intent-to-treat analysis of the DPP/DPPOS', *Diabetes Care*, vol. 35, no. 4, pp. 723–730, Apr. 2012, doi: 10.2337/dc11-1468.

[84]

Bailey, Stephen, *Academic Writing : A Handbook for International Students*, 5th edn. [Online]. Available: <https://ebookcentral.proquest.com/lib/leicester/detail.action?docID=5178437>

[85]

P. Creme and M. R. Lea, Writing At University: A Guide For Students, 3rd ed. Maidenhead: McGraw-Hill Education, 2008. [Electronic resource]. Available: <https://ebookcentral.proquest.com/lib/leicester/detail.action?pq-origsite=primo&docID=345134>