

## MD7002 - Prevention, Screening & Early Detection in Diabetes

View Online



Prevention, Screening & Early Detection in Diabetes

---

Absetz, P. et al. (2009) 'Type 2 Diabetes Prevention in the Real World: Three-year results of the GOAL Lifestyle Implementation Trial', *Diabetes Care*, 32(8), pp. 1418-1420. Available at: <https://doi.org/10.2337/dc09-0039>.

Ackermann, R.T. et al. (2008) 'Translating the Diabetes Prevention Program into the Community', *American Journal of Preventive Medicine*, 35(4), pp. 357-363. Available at: <https://doi.org/10.1016/j.amepre.2008.06.035>.

Carter, P., Khunti, K. and Davies, M.J. (2012) 'Dietary Recommendations for the Prevention of Type 2 diabetes: What Are They Based on?', *Journal of Nutrition and Metabolism*, 2012, pp. 1-6. Available at: <https://doi.org/10.1155/2012/847202>.

Chiasson, J.-L. et al. (2002) 'Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial', *The Lancet*, 359(9323), pp. 2072-2077. Available at: [https://doi.org/10.1016/S0140-6736\(02\)08905-5](https://doi.org/10.1016/S0140-6736(02)08905-5).

Church, T.S. et al. (2004) 'Exercise Capacity and Body Composition as Predictors of Mortality Among Men With Diabetes', *Diabetes Care*, 27(1), pp. 83-88. Available at: <https://doi.org/10.2337/diacare.27.1.83>.

Colberg, S.R. et al. (2010) 'Exercise and Type 2 Diabetes: The American College of Sports Medicine and the American Diabetes Association: joint position statement executive summary', *Diabetes Care*, 33(12), pp. 2692-2696. Available at: <https://doi.org/10.2337/dc10-1548>.

Cordain, L. et al. (1998) 'Physical Activity, Energy Expenditure and Fitness: An Evolutionary Perspective', *International Journal of Sports Medicine*, 19(05), pp. 328-335. Available at: <https://doi.org/10.1055/s-2007-971926>.

DeFronzo, R.A. et al. (2011) 'Pioglitazone for Diabetes Prevention in Impaired Glucose Tolerance', *New England Journal of Medicine*, 364(12), pp. 1104-1115. Available at: <https://doi.org/10.1056/NEJMoa1010949>.

Dyson, P.A. et al. (2011) 'Diabetes UK evidence-based nutrition guidelines for the prevention and management of diabetes', *Diabetic Medicine*, 28(11), pp. 1282-1288. Available at: <https://doi.org/10.1111/j.1464-5491.2011.03371.x>.

'Effect of Nateglinide on the Incidence of Diabetes and Cardiovascular Events' (2010) *New England Journal of Medicine*, 362(16), pp. 1463-1476. Available at: <https://doi.org/10.1056/NEJMoa1001122>.

- 'Effect of Valsartan on the Incidence of Diabetes' (2010) *New England Journal of Medicine*, 363(8), pp. 792–793. Available at: <https://doi.org/10.1056/NEJMc1005899>.
- Frank W. Booth (2000) 'Waging war on modern chronic diseases: primary prevention through exercise biology', *Journal of Applied Physiology*, 88(2), pp. 774–787. Available at: <http://jap.physiology.org/content/88/2/774>.
- Gillies, C.L. et al. (2007) 'Pharmacological and lifestyle interventions to prevent or delay type 2 diabetes in people with impaired glucose tolerance: systematic review and meta-analysis', *BMJ*, 334(7588), pp. 299–299. Available at: <https://doi.org/10.1136/bmj.39063.689375.55>.
- Gray, L.J. et al. (2011) 'Defining Obesity Cut-Off Points for Migrant South Asians', *PLoS ONE*, 6(10). Available at: <https://doi.org/10.1371/journal.pone.0026464>.
- Greaves, C.J. et al. (2011) 'Systematic review of reviews of intervention components associated with increased effectiveness in dietary and physical activity interventions', *BMC Public Health*, 11(1). Available at: <https://doi.org/10.1186/1471-2458-11-119>.
- Hamman, R.F. et al. (2006) 'Effect of Weight Loss With Lifestyle Intervention on Risk of Diabetes', *Diabetes Care*, 29(9), pp. 2102–2107. Available at: <https://doi.org/10.2337/dc06-0560>.
- Laaksonen, D.E. et al. (2005) 'Physical Activity in the Prevention of Type 2 Diabetes: The Finnish Diabetes Prevention Study', *Diabetes*, 54(1), pp. 158–165. Available at: <https://doi.org/10.2337/diabetes.54.1.158>.
- Lindstrom, J. et al. (2003) 'The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity', *Diabetes Care*, 26(12), pp. 3230–3236. Available at: <https://doi.org/10.2337/diacare.26.12.3230>.
- Norris, S.L. et al. (1996) 'Long-term non-pharmacological weight loss interventions for adults with prediabetes', in *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd. Available at: <https://doi.org/10.1002/14651858.CD005270>.
- 'Nutrition Recommendations and Interventions for Diabetes: A position statement of the American Diabetes Association' (2008) *Diabetes Care*, 31(Supplement 1), pp. S61–S78. Available at: <https://doi.org/10.2337/dc08-S061>.
- Pan, X.-R. et al. (1997) 'Effects of Diet and Exercise in Preventing NIDDM in People With Impaired Glucose Tolerance: The Da Qing IGT and Diabetes Study', *Diabetes Care*, 20(4), pp. 537–544. Available at: <https://doi.org/10.2337/diacare.20.4.537>.
- 'Public Health England - Diabetes prevention programmes: evidence review' (2015). Public Health England. Available at: <https://www.gov.uk/government/publications/diabetes-prevention-programmes-evidence-review>.
- Ramachandran, A. et al. (2006) 'The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1)', *Diabetologia*, 49(2), pp. 289–297. Available at:

<https://doi.org/10.1007/s00125-005-0097-z>.

'Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin' (2002) *New England Journal of Medicine*, 346(6), pp. 393–403. Available at: <https://doi.org/10.1056/NEJMoa012512>.

Schwarz, P.E. et al. (2012) 'Nonpharmacological interventions for the prevention of type 2 diabetes mellitus', *Nature Reviews Endocrinology* [Preprint]. Available at: <https://doi.org/10.1038/nrendo.2011.232>.

Schwarz, P.E.H. et al. (2007) 'Development and Implementation of A European Guideline and Training Standards for Diabetes Prevention-The IMAGE project', *Journal of Public Health*, 15(5), pp. 353–360. Available at: <https://doi.org/10.1007/s10389-007-0145-x>.

Schwarz, P.E.H. (2011) 'Public Health Implications: Translation into Diabetes Prevention Initiatives—Four-Level Public Health Concept', *Medical Clinics of North America*, 95(2), pp. 397–407. Available at: <https://doi.org/10.1016/j.mcna.2010.11.008>.

'Screening for type 2 diabetes: literature review and economic modelling' (2007) *Clinical Governance: An International Journal*, 12(4). Available at: <https://doi.org/10.1108/cgij.2007.24812dae.003>.

Steven N Blair (2009) 'Physical inactivity: the biggest public health problem of the 21st century', *British Journal of Sports Medicine*, 43, pp. 1–2. Available at: <http://bjsm.bmj.com/content/43/1/1.full>.

TELFORD, RICHARD D. (no date) 'Low Physical Activity and Obesity: Causes of Chronic Disease or Simply Predictors?', *Medicine & Science in Sports & Exercise*, 39(8). Available at: <http://ezproxy.lib.le.ac.uk/login?url=http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&AN=00005768-200708000-00005&LSLINK=80&D=ovft>.

The relative importance of modifiable potential risk factors of type 2 diabetes: a meta-analysis of two cohorts (no date). Available at: <http://link.springer.com/article/10.1007/s10654-009-9405-0/fulltext.html>.

Tuomilehto, J. et al. (2001) 'Prevention of Type 2 Diabetes Mellitus by Changes in Lifestyle among Subjects with Impaired Glucose Tolerance', *New England Journal of Medicine*, 344(18), pp. 1343–1350. Available at: <https://doi.org/10.1056/NEJM200105033441801>.

Umpierre, D. (2011) 'Physical Activity Advice Only or Structured Exercise Training and Association With HbA<sub>1c</sub> Levels in Type 2 Diabetes', *JAMA*, 305(17). Available at: <https://doi.org/10.1001/jama.2011.576>.

Yates, T. et al. (2009) 'Effectiveness of a Pragmatic Education Program Designed to Promote Walking Activity in Individuals With Impaired Glucose Tolerance: A randomized controlled trial', *Diabetes Care*, 32(8), pp. 1404–1410. Available at: <https://doi.org/10.2337/dc09-0130>.

Yates, T et al. (2009) 'The role of physical activity in the management of type 2 diabetes mellitus', *Postgraduate Medical Journal*, 85(1001), pp. 129–133. Available at:

<https://doi.org/10.1136/pgmj.2008.067900>.

Yates, T. et al. (2011) 'Stand up for your health: Is it time to rethink the physical activity paradigm?', *Diabetes Research and Clinical Practice*, 93(2), pp. 292–294. Available at: <https://doi.org/10.1016/j.diabres.2011.03.023>.