

MD7002 - Prevention, Screening & Early Detection in Diabetes

[View Online](#)

Prevention, Screening & Early Detection in Diabetes

[1]

'Public Health England - Diabetes prevention programmes: evidence review'. Public Health England, 2015. [Online]. Available: <https://www.gov.uk/government/publications/diabetes-prevention-programmes-evidence-review>

[2]

'Nutrition Recommendations and Interventions for Diabetes: A position statement of the American Diabetes Association', Diabetes Care, vol. 31, no. Supplement 1, pp. S61-S78, Jan. 2008, doi: 10.2337/dc08-S061.

[3]

P. Carter, K. Khunti, and M. J. Davies, 'Dietary Recommendations for the Prevention of Type 2 diabetes: What Are They Based on?', Journal of Nutrition and Metabolism, vol. 2012, pp. 1-6, 2012, doi: 10.1155/2012/847202.

[4]

S. R. Colberg et al., 'Exercise and Type 2 Diabetes: The American College of Sports Medicine and the American Diabetes Association: joint position statement executive summary', Diabetes Care, vol. 33, no. 12, pp. 2692-2696, Dec. 2010, doi: 10.2337/dc10-1548.

[5]

P. A. Dyson et al., 'Diabetes UK evidence-based nutrition guidelines for the prevention and management of diabetes', Diabetic Medicine, vol. 28, no. 11, pp. 1282-1288, Nov. 2011,

doi: 10.1111/j.1464-5491.2011.03371.x.

[6]

C. L. Gillies et al., 'Pharmacological and lifestyle interventions to prevent or delay type 2 diabetes in people with impaired glucose tolerance: systematic review and meta-analysis', *BMJ*, vol. 334, no. 7588, pp. 299–299, Jan. 2007, doi: 10.1136/bmj.39063.689375.55.

[7]

'Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin', *New England Journal of Medicine*, vol. 346, no. 6, pp. 393–403, Feb. 2002, doi: 10.1056/NEJMoa012512.

[8]

'The relative importance of modifiable potential risk factors of type 2 diabetes: a meta-analysis of two cohorts'. [Online]. Available: <http://link.springer.com/article/10.1007/s10654-009-9405-0/fulltext.html>

[9]

J. Lindstrom et al., 'The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity', *Diabetes Care*, vol. 26, no. 12, pp. 3230–3236, Dec. 2003, doi: 10.2337/diacare.26.12.3230.

[10]

X.-R. Pan et al., 'Effects of Diet and Exercise in Preventing NIDDM in People With Impaired Glucose Tolerance: The Da Qing IGT and Diabetes Study', *Diabetes Care*, vol. 20, no. 4, pp. 537–544, Apr. 1997, doi: 10.2337/diacare.20.4.537.

[11]

A. Ramachandran, C. Snehalatha, S. Mary, B. Mukesh, A. D. Bhaskar, and V. Vijay, 'The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1)', *Diabetologia*, vol. 49, no. 2, pp. 289–297, Feb. 2006, doi: 10.1007/s00125-005-0097-z.

[12]

P. E. Schwarz, C. J. Greaves, J. Lindström, T. Yates, and M. J. Davies, 'Nonpharmacological interventions for the prevention of type 2 diabetes mellitus', *Nature Reviews Endocrinology*, Jan. 2012, doi: 10.1038/nrendo.2011.232.

[13]

P. E. H. Schwarz, 'Public Health Implications: Translation into Diabetes Prevention Initiatives' Four-Level Public Health Concept, *Medical Clinics of North America*, vol. 95, no. 2, pp. 397–407, Mar. 2011, doi: 10.1016/j.mcna.2010.11.008.

[14]

J. Tuomilehto et al., 'Prevention of Type 2 Diabetes Mellitus by Changes in Lifestyle among Subjects with Impaired Glucose Tolerance', *New England Journal of Medicine*, vol. 344, no. 18, pp. 1343–1350, May 2001, doi: 10.1056/NEJM200105033441801.

[15]

'Screening for type 2 diabetes: literature review and economic modelling', *Clinical Governance: An International Journal*, vol. 12, no. 4, Oct. 2007, doi: 10.1108/cgij.2007.24812dae.003.

[16]

P. Absetz et al., 'Type 2 Diabetes Prevention in the Real World: Three-year results of the GOAL Lifestyle Implementation Trial', *Diabetes Care*, vol. 32, no. 8, pp. 1418–1420, Aug. 2009, doi: 10.2337/dc09-0039.

[17]

R. T. Ackermann, E. A. Finch, E. Brizendine, H. Zhou, and D. G. Marrero, 'Translating the Diabetes Prevention Program into the Community', *American Journal of Preventive Medicine*, vol. 35, no. 4, pp. 357–363, Oct. 2008, doi: 10.1016/j.amepre.2008.06.035.

[18]

J.-L. Chiasson, R. G. Josse, R. Gomis, M. Hanefeld, A. Karasik, and M. Laakso, 'Acarbose for prevention of type 2 diabetes mellitus: the STOP-NIDDM randomised trial', *The Lancet*, vol. 359, no. 9323, pp. 2072–2077, Jun. 2002, doi: 10.1016/S0140-6736(02)08905-5.

[19]

R. A. DeFronzo et al., 'Pioglitazone for Diabetes Prevention in Impaired Glucose Tolerance', *New England Journal of Medicine*, vol. 364, no. 12, pp. 1104–1115, Mar. 2011, doi: 10.1056/NEJMoa1010949.

[20]

L. J. Gray et al., 'Defining Obesity Cut-Off Points for Migrant South Asians', *PLoS ONE*, vol. 6, no. 10, Oct. 2011, doi: 10.1371/journal.pone.0026464.

[21]

C. J. Greaves et al., 'Systematic review of reviews of intervention components associated with increased effectiveness in dietary and physical activity interventions', *BMC Public Health*, vol. 11, no. 1, 2011, doi: 10.1186/1471-2458-11-119.

[22]

'Effect of Valsartan on the Incidence of Diabetes', *New England Journal of Medicine*, vol. 363, no. 8, pp. 792–793, Aug. 2010, doi: 10.1056/NEJMc1005899.

[23]

P. E. H. Schwarz, U. Gruhl, S. R. Bornstein, R. Landgraf, M. Hall, and J. Tuomilehto, 'Development and Implementation of A European Guideline and Training Standards for Diabetes Prevention-The IMAGE project', *Journal of Public Health*, vol. 15, no. 5, pp. 353–360, Sep. 2007, doi: 10.1007/s10389-007-0145-x.

[24]

'Effect of Nateglinide on the Incidence of Diabetes and Cardiovascular Events', *New England Journal of Medicine*, vol. 362, no. 16, pp. 1463–1476, Apr. 2010, doi: 10.1056/NEJMoa1001122.

[25]

T. Yates, M. Davies, T. Gorely, F. Bull, and K. Khunti, 'Effectiveness of a Pragmatic Education Program Designed to Promote Walking Activity in Individuals With Impaired Glucose Tolerance: A randomized controlled trial', *Diabetes Care*, vol. 32, no. 8, pp. 1404–1410, Aug. 2009, doi: 10.2337/dc09-0130.

[26]

T. Yates, E. G. Wilmot, K. Khunti, S. Biddle, T. Gorely, and M. J. Davies, 'Stand up for your health: Is it time to rethink the physical activity paradigm?', *Diabetes Research and Clinical Practice*, vol. 93, no. 2, pp. 292–294, Aug. 2011, doi: 10.1016/j.diabres.2011.03.023.

[27]

Steven N Blair, 'Physical inactivity: the biggest public health problem of the 21st century', *British Journal of Sports Medicine*, vol. 43, pp. 1–2, Jan. 2009, [Online]. Available: <http://bjsm.bmj.com/content/43/1/1.full>

[28]

Frank W. Booth, 'Waging war on modern chronic diseases: primary prevention through exercise biology', *Journal of Applied Physiology*, vol. 88, no. 2, pp. 774–787, 2000, [Online]. Available: <http://jap.physiology.org/content/88/2/774>

[29]

T. S. Church et al., 'Exercise Capacity and Body Composition as Predictors of Mortality Among Men With Diabetes', *Diabetes Care*, vol. 27, no. 1, pp. 83–88, Jan. 2004, doi: 10.2337/diacare.27.1.83.

[30]

L. Cordain, R. Gotshall, S. Eaton, and S. Eaton, 'Physical Activity, Energy Expenditure and Fitness: An Evolutionary Perspective', *International Journal of Sports Medicine*, vol. 19, no. 05, pp. 328–335, Jul. 1998, doi: 10.1055/s-2007-971926.

[31]

R. F. Hamman et al., 'Effect of Weight Loss With Lifestyle Intervention on Risk of Diabetes', *Diabetes Care*, vol. 29, no. 9, pp. 2102–2107, Sep. 2006, doi: 10.2337/dc06-0560.

[32]

D. E. Laaksonen et al., 'Physical Activity in the Prevention of Type 2 Diabetes: The Finnish Diabetes Prevention Study', *Diabetes*, vol. 54, no. 1, pp. 158–165, Jan. 2005, doi: 10.2337/diabetes.54.1.158.

[33]

S. L. Norris, X. Zhang, A. Avenell, E. Gregg, C. H. Schmid, and J. Lau, 'Long-term non-pharmacological weight loss interventions for adults with prediabetes', in *Cochrane Database of Systematic Reviews*, Chichester, UK: John Wiley & Sons, Ltd, 1996. doi: 10.1002/14651858.CD005270.

[34]

TELFORD, RICHARD D., 'Low Physical Activity and Obesity: Causes of Chronic Disease or Simply Predictors?', *Medicine & Science in Sports & Exercise*, vol. 39, no. 8, [Online]. Available: <http://ezproxy.lib.le.ac.uk/login?url=http://ovidsp.ovid.com/ovidweb.cgi?T=JS&CSC=Y&NEWS=N&PAGE=fulltext&AN=00005768-200708000-00005&LSLINK=80&D=ovft>

[35]

D. Umpierre, 'Physical Activity Advice Only or Structured Exercise Training and Association With HbA_{1c} Levels in Type 2 Diabetes', *JAMA*, vol. 305, no. 17, May 2011, doi: 10.1001/jama.2011.576.

[36]

T. Yates, K. Khunti, J. Troughton, and M. Davies, 'The role of physical activity in the management of type 2 diabetes mellitus', *Postgraduate Medical Journal*, vol. 85, no. 1001, pp. 129–133, Mar. 2009, doi: 10.1136/pgmj.2008.067900.