

MD7511 Injectable Therapies

Injectable Therapies

View Online



: 'Summary of Revisions' (2016) *Diabetes Care*, 39(Supplement 1), pp. S4–S5. Available at: <https://doi.org/10.2337/dc16-S003>.

A randomised, 52-week, treat-to-target trial comparing insulin detemir with insulin glargine when administered as add-on to glucose-lowering drugs in insulin-naive people with type 2 diabetes (no date). Available at: <http://link.springer.com.ezproxy4.lib.le.ac.uk/article/10.1007/s00125-007-0911-x/fulltext.html>.

Akram, J., De Verga, V. and Group, the R.S. (1999) 'Insulin lispro (Lys(B28), Pro(B29)) in the treatment of diabetes during the fasting month of Ramadan', *Diabetic Medicine*, 16(10), pp. 867–874. Available at: <https://doi.org/10.1046/j.1464-5491.1999.00164.x>.

Aldam, P., Levy, N. and Hall, G.M. (2014) 'Perioperative management of diabetic patients: new controversies', *British Journal of Anaesthesia*, 113(6), pp. 906–909. Available at: <https://doi.org/10.1093/bja/aeu259>.

Ali, S. et al. (2016) 'Guidelines for managing diabetes in Ramadan', *Diabetic Medicine*, 33(10), pp. 1315–1329. Available at: <https://doi.org/10.1111/dme.13080>.

Anderson, B., Funnell, M., and American Diabetes Association (2005) *The art of empowerment: stories and strategies for diabetes educators*. 2nd ed. Alexandria, Va: American Diabetes Association.

Anderson, J.W., Kendall, C.W.C. and Jenkins, D.J.A. (2003) 'Importance of Weight Management in Type 2 Diabetes: Review with Meta-analysis of Clinical Studies', *Journal of the American College of Nutrition*, 22(5), pp. 331–339. Available at: <https://doi.org/10.1080/07315724.2003.10719316>.

Ashwell, S.G. et al. (2006) 'Improved glycaemic control with insulin glargine plus insulin lispro: a multicentre, randomized, cross-over trial in people with Type 1 diabetes', *Diabetic Medicine*, 23(3), pp. 285–292. Available at: <https://doi.org/10.1111/j.1464-5491.2005.01781.x>.

Bailey, S. (2011) *Academic writing: a handbook for international students*. 3rd ed. London: Routledge. Available at: <http://ezproxy.lib.le.ac.uk/login?url=http://www.myilibrary.com?id=310467>.

Bajwa, S. et al. (2015) 'Interdisciplinary position statement on management of hyperglycemia in peri-operative and intensive care', *Journal of Anaesthesiology Clinical Pharmacology*, 31(2). Available at: <https://doi.org/10.4103/0970-9185.155141>.

Bakiner, O. et al. (2009) 'Repaglinide plus single-dose insulin glargine: a safe regimen for low-risk type 2 diabetic patients who insist on fasting in Ramadan', *Acta Diabetologica*, 46(1), pp. 63–65. Available at: <https://doi.org/10.1007/s00592-008-0062-7>.

Barker, P. et al. (2015) 'Peri-operative management of the surgical patient with diabetes 2015', *Anaesthesia*, 70(12), pp. 1427–1440. Available at: <https://doi.org/10.1111/anae.13233>.

'Basal Insulin and Cardiovascular and Other Outcomes in Dysglycemia' (2012) *New England Journal of Medicine*, 367(4), pp. 319–328. Available at: <https://doi.org/10.1056/NEJMoa1203858>.

Benkhadra, K. et al. (2016) 'Real Time Continuous Glucose Monitoring in type 1 diabetes: A Systematic review and Individual Patient Data Meta-Analysis', *Clinical Endocrinology* [Preprint]. Available at: <https://doi.org/10.1111/cen.13290>.

Bergenstal, R.M. et al. (2008) 'Adjust to Target in Type 2 Diabetes: Comparison of a simple algorithm with carbohydrate counting for adjustment of mealtime insulin glulisine', *Diabetes Care*, 31(7), pp. 1305–1310. Available at: <https://doi.org/10.2337/dc07-2137>.

Bergenstal, R.M. et al. (2010) 'Effectiveness of Sensor-Augmented Insulin-Pump Therapy in Type 1 Diabetes', *New England Journal of Medicine*, 363(4), pp. 311–320. Available at: <https://doi.org/10.1056/NEJMoa1002853>.

Bergenstal, R.M. et al. (2012) 'A Randomized, Controlled Study of Once-Daily LY2605541, a Novel Long-Acting Basal Insulin, Versus Insulin Glargine in Basal Insulin-Treated Patients With Type 2 Diabetes', *Diabetes Care*, 35(11), pp. 2140–2147. Available at: <https://doi.org/10.2337/dc12-0060>.

Bretzel, R.G. et al. (2008) 'Once-daily basal insulin glargine versus thrice-daily prandial insulin lispro in people with type 2 diabetes on oral hypoglycaemic agents (APOLLO): an open randomised controlled trial', *The Lancet*, 371(9618), pp. 1073–1084. Available at: [https://doi.org/10.1016/S0140-6736\(08\)60485-7](https://doi.org/10.1016/S0140-6736(08)60485-7).

Burnett, J.C.D. (2006) 'Long- and Short-Haul Travel by Air: Issues for People With Diabetes on Insulin', *Journal of Travel Medicine*, 13(5), pp. 255–260. Available at: <https://doi.org/10.1111/j.1708-8305.2006.00057.x>.

Buse, J.B. et al. (2011) 'Use of Twice-Daily Exenatide in Basal Insulin-Treated Patients With Type 2 Diabetes', *Annals of Internal Medicine*, 154(2). Available at: <https://doi.org/10.7326/0003-4819-154-2-201101180-00300>.

Buse, J.B. et al. (2014) 'Contribution of Liraglutide in the Fixed-Ratio Combination of Insulin Degludec and Liraglutide (IDegLira)', *Diabetes Care*, 37(11), pp. 2926–2933. Available at: <https://doi.org/10.2337/dc14-0785>.

'Clinical effectiveness and cost-effectiveness of continuous subcutaneous insulin infusion for diabetes: systematic review and economic evaluation' (no date). Available at: <http://www.journalslibrary.nihr.ac.uk/hta/volume-14/issue-11>.

Crasto, W. et al. (2009) 'Insulin U-500 in severe insulin resistance in type 2 diabetes mellitus', *Postgraduate Medical Journal*, 85(1002), pp. 219–222. Available at:

<https://doi.org/10.1136/pgmj.2008.073379>.

Crasto, W. (2016a) Handbook of Insulin Therapies. Adis; 1st ed. 2016 edition. Available at: <https://ebookcentral.proquest.com/lib/leicester/detail.action?docID=4676975>.

Crasto, W. (2016b) Handbook of Insulin Therapies. Adis; 1st ed. 2016 edition. Available at: <https://ebookcentral.proquest.com/lib/leicester/detail.action?docID=4676975>.

Crasto, W. (2016c) Handbook of Insulin Therapies. Adis; 1st ed. 2016 edition. Available at: <https://ebookcentral.proquest.com/lib/leicester/detail.action?docID=4676975>.

Crasto, W. (2016d) Handbook of Insulin Therapies. Adis; 1st ed. 2016 edition. Available at: <https://ebookcentral.proquest.com/lib/leicester/detail.action?docID=4676975>.

Crasto, W. (2016e) Handbook of Insulin Therapies. Adis; 1st ed. 2016 edition. Available at: <https://ebookcentral.proquest.com/lib/leicester/detail.action?docID=4676975>.

Crasto, W. (2016f) Handbook of Insulin Therapies. Adis; 1st ed. 2016 edition. Available at: <https://ebookcentral.proquest.com/lib/leicester/detail.action?docID=4676975>.

Daly, H. et al. (2015) 'Development of a self-management education module for those with type 2 diabetes on injectable therapies', *Practical Diabetes*, 32(8), pp. 305–310a. Available at: <https://doi.org/10.1002/pdi.1979>.

Davies, M. et al. (2005a) 'Improvement of Glycemic Control in Subjects With Poorly Controlled Type 2 Diabetes: Comparison of two treatment algorithms using insulin glargine', *Diabetes Care*, 28(6), pp. 1282–1288. Available at: <https://doi.org/10.2337/diacare.28.6.1282>.

Davies, M. et al. (2005b) 'Improvement of Glycemic Control in Subjects With Poorly Controlled Type 2 Diabetes: Comparison of two treatment algorithms using insulin glargine', *Diabetes Care*, 28(6), pp. 1282–1288. Available at: <https://doi.org/10.2337/diacare.28.6.1282>.

Davies, M.J. et al. (2009) 'Exenatide compared with long-acting insulin to achieve glycaemic control with minimal weight gain in patients with type 2 diabetes: results of the Helping Evaluate Exenatide in patients with diabetes compared with Long-Acting insulin (HEELA) study', *Diabetes, Obesity and Metabolism*, 11(12), pp. 1153–1162. Available at: <https://doi.org/10.1111/j.1463-1326.2009.01154.x>.

Davies, M.J. et al. (2014) 'Efficacy and safety of insulin degludec given as part of basal-bolus treatment with mealtime insulin aspart in type 1 diabetes: a 26-week randomized, open-label, treat-to-target non-inferiority trial', *Diabetes, Obesity and Metabolism*, 16(10), pp. 922–930. Available at: <https://doi.org/10.1111/dom.12298>.

Davies, M.J. et al. (2017) 'Impact of baseline HbA1c, diabetes duration and BMI on clinical outcomes in the LixiLan-O trial testing iGlarLixi (insulin glargine/lixisenatide titratable fixed-ratio combination) versus insulin glargine and lixisenatide monocomponents', *Diabetes, Obesity and Metabolism* [Preprint]. Available at: <https://doi.org/10.1111/dom.12980>.

Davies, M.J. and Chatterjee, S. (2017) 'Trial watch: Insulin initiation for type 2 diabetes

mellitus in primary care', *Nature Reviews Endocrinology*, 13(6), pp. 317–318. Available at: <https://doi.org/10.1038/nrendo.2017.41>.

Diabetes & Metabolism - Présentation - EM consulte (no date). Available at: <http://www.em-consulte.com/article/80031/alertePM#N1010C>.

Diamant, M. et al. (2010) 'Once weekly exenatide compared with insulin glargine titrated to target in patients with type 2 diabetes (DURATION-3): an open-label randomised trial', *The Lancet*, 375(9733), pp. 2234–2243. Available at: [https://doi.org/10.1016/S0140-6736\(10\)60406-0](https://doi.org/10.1016/S0140-6736(10)60406-0).

EASD Virtual Meeting (no date). Available at: <http://www.easdvirtualmeeting.org/resources/systematic-review-and-meta-analysis-of-the-efficacy-and-safety-of-sgl2-inhibitors-in-patients-with-type-2-diabetes-mellitus>.

'Effects of Intensive Glucose Lowering in Type 2 Diabetes' (2008) *New England Journal of Medicine*, 358(24), pp. 2545–2559. Available at: <https://doi.org/10.1056/NEJMoa0802743>.
Efficacy and safety of a fixed-ratio combination of insulin degludec and liraglutide (IDegLira) compared with its components given alone: results of a phase 3, open-label, randomised, 26-week, treat-to-target trial in insulin-naive patients with type 2 diabetes - ClinicalKey (no date). Available at: <https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/journal/1-s2.0-S2213858714701743>.

'Efficacy and Safety of Glucagon-like peptide-1 receptor agonists in type 2 diabetes Systematic review and mixed-treatment comparison analysis' (no date). Available at: <http://onlinelibrary.wiley.com.ezproxy3.lib.le.ac.uk/doi/10.1111/dom.12849/epdf>.

Eng, C. et al. (2014) 'Glucagon-like peptide-1 receptor agonist and basal insulin combination treatment for the management of type 2 diabetes: a systematic review and meta-analysis', *The Lancet*, 384(9961), pp. 2228–2234. Available at: [https://doi.org/10.1016/S0140-6736\(14\)61335-0](https://doi.org/10.1016/S0140-6736(14)61335-0).

Evans, M. et al. (2011) 'A review of modern insulin analogue pharmacokinetic and pharmacodynamic profiles in type 2 diabetes: improvements and limitations', *Diabetes, Obesity and Metabolism*, 13(8), pp. 677–684. Available at: <https://doi.org/10.1111/j.1463-1326.2011.01395.x>.

F Game (2012) 'Annals of The Royal College of Surgeons of England', 94(5). Available at: <https://doi.org/doi:10.1308/003588412X13171221591655>.

Fineman, M.S. et al. (2012) 'GLP-1 based therapies: differential effects on fasting and postprandial glucose', *Diabetes, Obesity and Metabolism*, 14(8), pp. 675–688. Available at: <https://doi.org/10.1111/j.1463-1326.2012.01560.x>.

Frandsen, C.S.S. and Madsbad, S. (2014a) 'Efficacy and safety of dipeptidyl peptidase-4 inhibitors as an add-on to insulin treatment in patients with Type 2 diabetes: a review', *Diabetic Medicine*, 31(11), pp. 1293–1300. Available at: <https://doi.org/10.1111/dme.12561>.

Frandsen, C.S.S. and Madsbad, S. (2014b) 'Efficacy and safety of dipeptidyl peptidase-4 inhibitors as an add-on to insulin treatment in patients with Type 2 diabetes: a review',

Diabetic Medicine, 31(11), pp. 1293–1300. Available at:
<https://doi.org/10.1111/dme.12561>.

Frisch, A. et al. (2010) 'Prevalence and Clinical Outcome of Hyperglycemia in the Perioperative Period in Noncardiac Surgery', *Diabetes Care*, 33(8), pp. 1783–1788. Available at: <https://doi.org/10.2337/dc10-0304>.

Fullerton, B. et al. (1996) 'Intensive glucose control versus conventional glucose control for type 1 diabetes mellitus', in *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd. Available at: <https://doi.org/10.1002/14651858.CD009122.pub2>.

Garber, A.J. et al. (2006) 'Attainment of glycaemic goals in type 2 diabetes with once-, twice-, or thrice-daily dosing with biphasic insulin aspart 70/30 (The 1-2-3 study)', *Diabetes, Obesity and Metabolism*, 8(1), pp. 58–66. Available at: <https://doi.org/10.1111/j.1463-1326.2005.00563.x>.

Garber, A.J. et al. (2007) 'Premixed insulin treatment for type 2 diabetes: analogue or human?', *Diabetes, Obesity and Metabolism*, 9(5), pp. 630–639. Available at: <https://doi.org/10.1111/j.1463-1326.2006.00654.x>.

Giugliano, D. et al. (2011) 'Efficacy of Insulin Analogs in Achieving the Hemoglobin A1c Target of <7% in Type 2 Diabetes: Meta-analysis of randomized controlled trials', *Diabetes Care*, 34(2), pp. 510–517. Available at: <https://doi.org/10.2337/dc10-1710>.

Goudswaard, A.N. et al. (1996) 'Insulin monotherapy versus combinations of insulin with oral hypoglycaemic agents in patients with type 2 diabetes mellitus', in *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd. Available at: <https://doi.org/10.1002/14651858.CD003418.pub2>.

Gough, S.C.L. et al. (2013) 'Low-Volume Insulin Degludec 200 Units/mL Once Daily Improves Glycemic Control Similarly to Insulin Glargine With a Low Risk of Hypoglycemia in Insulin-Naive Patients With Type 2 Diabetes: A 26-week, randomized, controlled, multinational, treat-to-target trial: The BEGIN LOW VOLUME trial', *Diabetes Care*, 36(9), pp. 2536–2542. Available at: <https://doi.org/10.2337/dc12-2329>.

Gregory T. Mucha, Stacia Merkel, William Thomas, John P. Bantle (no date) 'Fasting and insulin glargine in individuals with type 1 diabetes', *Diabetes Care* [Preprint]. Available at: http://go.galegroup.com.ezproxy4.lib.le.ac.uk/ps/retrieve.do?tabID=T002&resultListType=RESULT_LIST&searchResultsType=SingleTab&searchType=AdvancedSearchForm¤tPosition=17&docId=GALE%7CA116666633&docType=Article&sort=Relevance&contentSegment=&prodId=EAIM&contentSet=GALE%7CA116666633&searchId=R8&userGroupName=leicester&inPS=true.

Guevara-Aguirre, J. et al. (2004) 'Beneficial Effects of Addition of Oral Spray Insulin (Oralin) on Insulin Secretion and Metabolic Control in Subjects with Type 2 Diabetes Mellitus Suboptimally Controlled on Oral Hypoglycemic Agents', *Diabetes Technology & Therapeutics*, 6(1), pp. 1–8. Available at: <https://doi.org/10.1089/152091504322783341>.

Gururaj Setty, S. et al. (2016) 'New insulins and newer insulin regimens: a review of their role in improving glycaemic control in patients with diabetes', *Postgraduate Medical Journal*, 92(1085), pp. 152–164. Available at: <https://doi.org/10.1136/postgradmedj-2015-133716>.

Gurwitz, J.H. (1994) 'Glucocorticoids and the Risk for Initiation of Hypoglycemic Therapy', *Archives of Internal Medicine*, 154(1). Available at: <https://doi.org/10.1001/archinte.1994.00420010131015>.

Hannele Yki-Järvinen (2013) 'Is There Evidence to Support Use of Premixed or Prandial Insulin Regimens in Insulin-Naive or Previously Insulin-Treated Type 2 Diabetic Patients?', *Diabetes Care*, 36(Suppl 2). Available at: <https://doi.org/doi:10.2337/dcS13-2026>.

Hannele Yki-Jarvinen, Leena Juurinen, Michael Alvarsson, Tord Bystedt, Ian Caldwell, Melanie Davies (no date) 'Initiate insulin by aggressive titration and education (Initiate): randomized study to compare initiation of insulin combination therapy in type 2 diabetic patients individually and in groups', *Diabetes Care* [Preprint]. Available at: http://go.galegroup.com.ezproxy4.lib.le.ac.uk/ps/retrieve.do?sort=DA-SORT&docType=Clinical+report&tabID=T002&prodId=EAIM&searchId=R1&resultListType=RESULT_LIST&searchType=AdvancedSearchForm&contentSegment=¤tPosition=5&searchResultsType=SingleTab&inPS=true&userGroupName=leicester&docId=GALE%7CA165017903&contentSet=GALE%7CA165017903.

Heller, S. et al. (2012) 'Insulin degludec, an ultra-longacting basal insulin, versus insulin glargine in basal-bolus treatment with mealtime insulin aspart in type 1 diabetes (BEGIN Basal-Bolus Type 1): a phase 3, randomised, open-label, treat-to-target non-inferiority trial', *The Lancet*, 379(9825), pp. 1489–1497. Available at: [https://doi.org/10.1016/S0140-6736\(12\)60204-9](https://doi.org/10.1016/S0140-6736(12)60204-9).

Heller, S.R. et al. (2004) 'Hypoglycaemia with insulin aspart: a double-blind, randomised, crossover trial in subjects with Type 1 diabetes', *Diabetic Medicine*, 21(7), pp. 769–775. Available at: <https://doi.org/10.1111/j.1464-5491.2004.01244.x>.

Hermansen, K. et al. (2006) 'A 26-Week, Randomized, Parallel, Treat-to-Target Trial Comparing Insulin Detemir With NPH Insulin as Add-On Therapy to Oral Glucose-Lowering Drugs in Insulin-Naive People With Type 2 Diabetes', *Diabetes Care*, 29(6), pp. 1269–1274. Available at: <https://doi.org/10.2337/dc05-1365>.

Hermansen, K. and Davies, M. (2007) 'Does insulin detemir have a role in reducing risk of insulin-associated weight gain?', *Diabetes, Obesity and Metabolism*, 9(3), pp. 209–217. Available at: <https://doi.org/10.1111/j.1463-1326.2006.00665.x>.

Hirsch, I.B. et al. (2014) 'Options for prandial glucose management in type 2 diabetes patients using basal insulin: addition of a short-acting GLP-1 analogue versus progression to basal-bolus therapy', *Diabetes, Obesity and Metabolism*, 16(3), pp. 206–214. Available at: <https://doi.org/10.1111/dom.12136>.

Hirsch, I.B. et al. (2017) 'Safety and efficacy of insulin degludec/insulin aspart with bolus mealtime insulin aspart compared with standard basal-bolus treatment in people with Type 1 diabetes: 1-year results from a randomized clinical trial (BOOST T1)', *Diabetic Medicine*, 34(2), pp. 167–173. Available at: <https://doi.org/10.1111/dme.13068>.

Holman, R.R. et al. (2007a) 'Addition of Biphasic, Prandial, or Basal Insulin to Oral Therapy in Type 2 Diabetes', *New England Journal of Medicine*, 357(17), pp. 1716–1730. Available at: <https://doi.org/10.1056/NEJMoa075392>.

- Holman, R.R. et al. (2007b) 'Addition of Biphasic, Prandial, or Basal Insulin to Oral Therapy in Type 2 Diabetes', *New England Journal of Medicine*, 357(17), pp. 1716–1730. Available at: <https://doi.org/10.1056/NEJMoa075392>.
- Holman, R.R. et al. (2008) '10-Year Follow-up of Intensive Glucose Control in Type 2 Diabetes', *New England Journal of Medicine*, 359(15), pp. 1577–1589. Available at: <https://doi.org/10.1056/NEJMoa0806470>.
- Holman, R.R. et al. (2009a) 'Three-Year Efficacy of Complex Insulin Regimens in Type 2 Diabetes', *New England Journal of Medicine*, 361(18), pp. 1736–1747. Available at: <https://doi.org/10.1056/NEJMoa0905479>.
- Holman, R.R. et al. (2009b) 'Three-Year Efficacy of Complex Insulin Regimens in Type 2 Diabetes', *New England Journal of Medicine*, 361(18), pp. 1736–1747. Available at: <https://doi.org/10.1056/NEJMoa0905479>.
- Holman, R.R. et al. (2009c) 'Three-Year Efficacy of Complex Insulin Regimens in Type 2 Diabetes', *New England Journal of Medicine*, 361(18), pp. 1736–1747. Available at: <https://doi.org/10.1056/NEJMoa0905479>.
- Holman, R.R. and Turner, R.C. (1985) 'A Practical Guide to Basal and Prandial Insulin Therapy', *Diabetic Medicine*, 2(1), pp. 45–53. Available at: <https://doi.org/10.1111/j.1464-5491.1985.tb00592.x>.
- Holt, R.I.G. (2010a) *Textbook of diabetes*. 4th ed. Chichester: Wiley-Blackwell. Available at: <http://ezproxy.lib.le.ac.uk/login?url=http://www.myilibrary.com?id=269077>.
- Holt, R.I.G. (2010b) *Textbook of diabetes*. 4th ed. Chichester: Wiley-Blackwell. Available at: <http://ezproxy.lib.le.ac.uk/login?url=http://www.myilibrary.com?id=269077>.
- Holt, R.I.G. et al. (2016a) *Textbook of Diabetes*. 5th ed. Somerset: John Wiley & Sons, Incorporated. Available at: <http://ebookcentral.proquest.com/lib/leicester/detail.action?docID=4769056>.
- Holt, R.I.G. et al. (2016b) *Textbook of Diabetes*. 5th ed. Somerset: John Wiley & Sons, Incorporated. Available at: <http://ebookcentral.proquest.com/lib/leicester/detail.action?docID=4769056>.
- Home, P. et al. (2014) 'Insulin Therapy in People With Type 2 Diabetes: Opportunities and Challenges?', *Diabetes Care*, 37(6), pp. 1499–1508. Available at: <https://doi.org/10.2337/dc13-2743>.
- Home, P.D. (2012) 'The pharmacokinetics and pharmacodynamics of rapid-acting insulin analogues and their clinical consequences', *Diabetes, Obesity and Metabolism*, 14(9), pp. 780–788. Available at: <https://doi.org/10.1111/j.1463-1326.2012.01580.x>.
- Home, P.D. et al. (2015) 'New Insulin Glargine 300 Units/mL Versus Glargine 100 Units/mL in People With Type 1 Diabetes: A Randomized, Phase 3a, Open-Label Clinical Trial (EDITION 4)', *Diabetes Care*, 38(12), pp. 2217–2225. Available at: <https://doi.org/10.2337/dc15-0249>.
- Hommel, I. et al. (2015) 'Perioperative diabetes care: room for improving the person

centredness', *Diabetic Medicine*, 32(4), pp. 561–568. Available at: <https://doi.org/10.1111/dme.12600>.

Horvath, K. et al. (1996a) 'Long-acting insulin analogues versus NPH insulin (human isophane insulin) for type 2 diabetes mellitus', in *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd. Available at: <https://doi.org/10.1002/14651858.CD005613.pub3>.

Horvath, K. et al. (1996b) 'Long-acting insulin analogues versus NPH insulin (human isophane insulin) for type 2 diabetes mellitus', in *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd. Available at: <https://doi.org/10.1002/14651858.CD005613.pub3>.

Hui, E. et al. (2010a) 'Comparison of Humalog Mix 50 with human insulin Mix 30 in type 2 diabetes patients during Ramadan', *International Journal of Clinical Practice*, 64(8), pp. 1095–1099. Available at: <https://doi.org/10.1111/j.1742-1241.2010.02347.x>.

Hui, E. et al. (2010b) 'Comparison of Humalog Mix 50 with human insulin Mix 30 in type 2 diabetes patients during Ramadan', *International Journal of Clinical Practice*, 64(8), pp. 1095–1099. Available at: <https://doi.org/10.1111/j.1742-1241.2010.02347.x>.

Hwang, J.L. and Weiss, R.E. (2014) 'Steroid-induced diabetes: a clinical and molecular approach to understanding and treatment', *Diabetes/Metabolism Research and Reviews*, 30(2), pp. 96–102. Available at: <https://doi.org/10.1002/dmrr.2486>.

'Hyperglycemia and Adverse Pregnancy Outcomes' (2008) *New England Journal of Medicine*, 358(19), pp. 1991–2002. Available at: <https://doi.org/10.1056/NEJMoa0707943>.
Insulin glargine or NPH combined with metformin in type 2 diabetes: the LANMET study (no date). Available at: <http://link.springer.com.ezproxy4.lib.le.ac.uk/article/10.1007/s00125-005-0132-0/fulltext.html>.

Insulin treatment and the problem of weight gain in type 2 diabetes. - PubMed - NCBI (no date). Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17102158>.

Intensity of peri-operative glycemic control and postoperative outcomes in patients with diabetes: a meta-analysis - ClinicalKey (no date). Available at: <https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S0168822713001873>.

Ismail-Beigi, F. (2012) 'Glycemic Management of Type 2 Diabetes Mellitus', *New England Journal of Medicine*, 366(14), pp. 1319–1327. Available at: <https://doi.org/10.1056/NEJMcp1013127>.

Jackson, M.J. et al. (2016) 'Perioperative management of diabetes in elective patients: a region-wide audit', *British Journal of Anaesthesia*, 116(4), pp. 501–506. Available at: <https://doi.org/10.1093/bja/aev554>.

Jenkins, K. (2014) 'II. Needle phobia: a psychological perspective', *British Journal of Anaesthesia*, 113(1), pp. 4–6. Available at: <https://doi.org/10.1093/bja/aeu013>.

June James (no date) 'End-of-life care: Anything but a pathway'. Available at:

http://www.diabetesandprimarycare.co.uk/media/content/_master/3572/files/pdf/dpc15-6-292-7.pdf.

June James (no date) 'Safety and insulin: implementation of national guidance at a local level', *Journal of Diabetes Nursing* [Preprint]. Available at: http://go.galegroup.com.ezproxy4.lib.le.ac.uk/ps/retrieve.do?tabID=T002&resultListType=RESULT_LIST&searchResultsType=SingleTab&searchType=AdvancedSearchForm¤tPosition=8&docId=GALE%7CA342482979&docType=Article&sort=RELEVANCE&contentSegment=&prodId=EAIM&contentSet=GALE%7CA342482979&searchId=R6&userGroupName=leicester&inPS=true.

Kahler, P. et al. (2014) 'Targeting intensive versus conventional glycaemic control for type 1 diabetes mellitus: a systematic review with meta-analyses and trial sequential analyses of randomised clinical trials', *BMJ Open*, 4(8), pp. e004806–e004806. Available at: <https://doi.org/10.1136/bmjopen-2014-004806>.

Khedkar, A. et al. (2010) 'A dose range finding study of novel oral insulin (IN-105) under fed conditions in type 2 diabetes mellitus subjects', *Diabetes, Obesity and Metabolism*, 12(8), pp. 659–664. Available at: <https://doi.org/10.1111/j.1463-1326.2010.01213.x>.

Kramer, C.K., Zinman, B. and Retnakaran, R. (2013) 'Short-term intensive insulin therapy in type 2 diabetes mellitus: a systematic review and meta-analysis', *The Lancet Diabetes & Endocrinology*, 1(1), pp. 28–34. Available at: [https://doi.org/10.1016/S2213-8587\(13\)70006-8](https://doi.org/10.1016/S2213-8587(13)70006-8).

Kuhlmann, M. and Marre, M. (2010) 'Lessons learned from biosimilar epoetins and insulins', *The British Journal of Diabetes & Vascular Disease*, 10(2), pp. 90–97. Available at: <https://doi.org/10.1177/1474651409355454>.

Kutz, A. et al. (2017) 'The association of admission hyperglycaemia and adverse clinical outcome in medical emergencies: the multinational, prospective, observational TRIAGE study', *Diabetic Medicine* [Preprint]. Available at: <https://doi.org/10.1111/dme.13325>.

L. Luzi (1989) 'Effect of loss of first-phase insulin secretion on hepatic glucose production and tissue glucose disposal in humans', *American Journal of Physiology - Endocrinology and Metabolism*, 257(2), pp. E241–E246. Available at: <http://ajpendo.physiology.org.ezproxy4.lib.le.ac.uk/content/257/2/E241>.

Lambert, K. and Holt, R.I.G. (2013) 'The use of insulin analogues in pregnancy', *Diabetes, Obesity and Metabolism*, 15(10), pp. 888–900. Available at: <https://doi.org/10.1111/dom.12098>.

Lankisch, M.R. et al. (2008) 'Introducing a simplified approach to insulin therapy in type 2 diabetes: a comparison of two single-dose regimens of insulin glulisine plus insulin glargine and oral antidiabetic drugs', *Diabetes, Obesity and Metabolism* [Preprint]. Available at: <https://doi.org/10.1111/j.1463-1326.2008.00967.x>.

Lasserson, D.S. et al. (2009a) 'Optimal insulin regimens in type 2 diabetes mellitus: systematic review and meta-analyses', *Diabetologia*, 52(10), pp. 1990–2000. Available at: <https://doi.org/10.1007/s00125-009-1468-7>.

Lasserson, D.S. et al. (2009b) 'Optimal insulin regimens in type 2 diabetes mellitus:

systematic review and meta-analyses', *Diabetologia*, 52(10), pp. 1990–2000. Available at: <https://doi.org/10.1007/s00125-009-1468-7>.

Lea, Mary R. □Creme, Phyllis (2008) *Writing At University*. 3rd edn. McGraw-Hill Education. Available at: <http://ebookcentral.proquest.com.ezproxy3.lib.le.ac.uk/lib/leicester/detail.action?docID=345134>.

Levy, N., Penfold, N. and Mythen, M. (2016) 'Perioperative management of diabetes and the emerging role of anaesthetists as perioperative physicians', *British Journal of Anaesthesia*, 116(4), pp. 443–447. Available at: <https://doi.org/10.1093/bja/aew049>.

Liu, D. et al. (2013) 'A practical guide to the monitoring and management of the complications of systemic corticosteroid therapy', *Allergy, Asthma & Clinical Immunology*, 9(1). Available at: <https://doi.org/10.1186/1710-1492-9-30>.

Luc JC van Loon (2000) 'Plasma insulin responses after ingestion of different amino acid or protein mixtures with carbohydrate', *The American Journal of Clinical Nutrition*, 72(1), pp. 96–105. Available at: <http://ajcn.nutrition.org/content/72/1/96.full>.

Luzio, S.D. et al. (2010) 'The glucose lowering effect of an oral insulin (Capsulin) during an isoglycaemic clamp study in persons with type 2 diabetes', *Diabetes, Obesity and Metabolism*, 12(1), pp. 82–87. Available at: <https://doi.org/10.1111/j.1463-1326.2009.01146.x>.

Manju Chandran, Steven V. Edelman (no date) 'Have insulin, will fly: diabetes management during air travel and time zone adjustment strategies. (Practical Pointers)', *Clinical Diabetes* [Preprint]. Available at: http://go.galegroup.com.ezproxy3.lib.le.ac.uk/ps/retrieve.do?tabID=T002&resultListType=RESULT_LIST&searchResultsType=SingleTab&searchType=AdvancedSearchForm¤tPosition=3&docId=GALE%7CA100730300&docType=Article&sort=Relevance&contentSegment=&prodId=EAIM&contentSet=GALE%7CA100730300&searchId=R2&userGroupName=leicester&inPS=true.

Martha M. Funnell (no date) 'Overcoming barriers to the initiation of insulin therapy', *Clinical Diabetes* [Preprint]. Available at: http://go.galegroup.com.ezproxy3.lib.le.ac.uk/ps/retrieve.do?sort=DA-SORT&docType=Article&tabID=T002&prodId=EAIM&searchId=R1&resultListType=RESULT_LIST&searchType=AdvancedSearchForm&contentSegment=¤tPosition=8&searchResultsType=SingleTab&inPS=true&userGroupName=leicester&docId=GALE%7CA159389875&contentSet=GALE%7CA159389875.

Matthew C. Riddle, Julio Rosenstock, John Gerich (no date) 'The treat-to-target trial: randomized addition of glargine or human NPH insulin to oral therapy of type 2 diabetic patients', *Diabetes Care* [Preprint]. Available at: http://go.galegroup.com.ezproxy4.lib.le.ac.uk/ps/retrieve.do?tabID=T002&resultListType=RESULT_LIST&searchResultsType=SingleTab&searchType=AdvancedSearchForm¤tPosition=9&docId=GALE%7CA110470954&docType=Article&sort=Relevance&contentSegment=&prodId=EAIM&contentSet=GALE%7CA110470954&searchId=R3&userGroupName=leicester&inPS=true.

Mattoo, V. et al. (2003) 'A comparison of insulin lispro Mix25™ and human insulin 30/70 in

the treatment of type 2 diabetes during Ramadan', *Diabetes Research and Clinical Practice*, 59(2), pp. 137–143. Available at: [https://doi.org/10.1016/S0168-8227\(02\)00202-4](https://doi.org/10.1016/S0168-8227(02)00202-4).

McBain, H. et al. (2017) 'Barriers to and enablers of insulin self-titration in adults with Type 2 diabetes: a qualitative study', *Diabetic Medicine*, 34(2), pp. 253–261. Available at: <https://doi.org/10.1111/dme.13196>.

McCrimmon, R.J. (2017) 'Old habits are hard to break: lessons from the study of hypoglycaemia', *Diabetic Medicine*, 34(2), pp. 148–155. Available at: <https://doi.org/10.1111/dme.13277>.

Menting, J.G. et al. (2013) 'How insulin engages its primary binding site on the insulin receptor', *Nature*, 493(7431), pp. 241–245. Available at: <https://doi.org/10.1038/nature11781>.

Mollema, E.D. et al. (2001) 'Phobia of self-injecting and self-testing in insulin-treated diabetes patients: opportunities for screening', *Diabetic Medicine*, 18(8), pp. 671–674. Available at: <https://doi.org/10.1046/j.1464-5491.2001.00547.x>.

Morçöl, T. et al. (2004) 'Calcium phosphate-PEG-insulin-casein (CAPIC) particles as oral delivery systems for insulin', *International Journal of Pharmaceutics*, 277(1–2), pp. 91–97. Available at: <https://doi.org/10.1016/j.ijpharm.2003.07.015>.

Nathan, D.M. et al. (2006) 'Management of Hyperglycemia in Type 2 Diabetes: A Consensus Algorithm for the Initiation and Adjustment of Therapy: A consensus statement from the American Diabetes Association and the European Association for the Study of Diabetes', *Diabetes Care*, 29(8), pp. 1963–1972. Available at: <https://doi.org/10.2337/dc06-9912>.

Nathan, D.M. (2014) 'The Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications Study at 30 Years: Overview', *Diabetes Care*, 37(1), pp. 9–16. Available at: <https://doi.org/10.2337/dc13-2112>.

Nauck, M. (2016) 'Incretin therapies: highlighting common features and differences in the modes of action of glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors', *Diabetes, Obesity and Metabolism*, 18(3), pp. 203–216. Available at: <https://doi.org/10.1111/dom.12591>.

New IDegLira data show rapid and predictable glycaemic improvements in people with type 2 diabetes (no date). Available at: <http://www.multivu.com/players/English/72762519-novo-nordisk-IDegLira-treatment/>.

New Insulin Glargine 300 U/mL: Glycemic Control and Hypoglycemia in a Meta-Analysis of Phase 3a EDITION Clinical Trials in People with T2DM - ClinicalKey (no date). Available at: <https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S1499267114003086>.

Owens, D.R. et al. (2011) 'Effects of initiation and titration of a single pre-prandial dose of insulin glulisine while continuing titrated insulin glargine in type 2 diabetes: a 6-month 'proof-of-concept' study', *Diabetes, Obesity and Metabolism*, 13(11), pp. 1020–1027. Available at: <https://doi.org/10.1111/j.1463-1326.2011.01459.x>.

Owens, D.R. (2013) 'Stepwise intensification of insulin therapy in Type 2 diabetes management-exploring the concept of the basal-plus approach in clinical practice', *Diabetic Medicine*, 30(3), pp. 276–288. Available at: <https://doi.org/10.1111/dme.12019>.

P. Schneiter (1998) 'Kinetics of dexamethasone-induced alterations of glucose metabolism in healthy humans', *American Journal of Physiology - Endocrinology and Metabolism*, 275(5), pp. E806–E813. Available at: <http://ajpendo.physiology.org/content/275/5/E806>.

Park, S.W. et al. (2017) 'Insulin degludec/insulin aspart once daily in Type 2 diabetes: a comparison of simple or stepwise titration algorithms (BOOST : SIMPLE USE)', *Diabetic Medicine*, 34(2), pp. 174–179. Available at: <https://doi.org/10.1111/dme.13069>.

Peter Mansell (no date) 'The dose adjustment for normal eating (DAFNE) education programme', *Journal of Diabetes Nursing* [Preprint]. Available at: http://go.galegroup.com.ezproxy4.lib.le.ac.uk/ps/retrieve.do?tabID=T002&resultListType=RESULT_LIST&searchResultsType=SingleTab&searchType=AdvancedSearchForm¤tPosition=1&docId=GALE%7CA314268073&docType=Article&sort=Relevance&contentSegment=&prodId=EAIM&contentSet=GALE%7CA314268073&searchId=R1&userGroupName=leicester&inPS=true.

Philip Raskin, Elsie Allen, Priscilla Hollander, Andrew Lewin, Robert A. Gabbay, Peter Hu (no date) 'Initiating insulin therapy in type 2 diabetes: a comparison of biphasic and basal insulin analogs', *Diabetes Care* [Preprint]. Available at: http://go.galegroup.com.ezproxy4.lib.le.ac.uk/ps/retrieve.do?sort=DA-SORT&docType=Article&tabID=T002&prodId=EAIM&searchId=R5&resultListType=RESULT_LIST&searchType=AdvancedSearchForm&contentSegment=¤tPosition=5&searchResultsType=SingleTab&inPS=true&userGroupName=leicester&docId=GALE%7CA128867472&contentSet=GALE%7CA128867472.

Pickup, J.C., Reznik, Y. and Sutton, A.J. (2017) 'Glycemic Control During Continuous Subcutaneous Insulin Infusion Versus Multiple Daily Insulin Injections in Type 2 Diabetes: Individual Patient Data Meta-analysis and Meta-regression of Randomized Controlled Trials', *Diabetes Care*, 40(5), pp. 715–722. Available at: <https://doi.org/10.2337/dc16-2201>.

Pozzilli, P. et al. (2005) 'Biokinetics of buccal spray insulin in patients with type 1 diabetes', *Metabolism*, 54(7), pp. 930–934. Available at: <https://doi.org/10.1016/j.metabol.2005.02.008>.

Prandial inhaled insulin plus basal insulin glargine versus twice daily biaspart insulin for type 2 diabetes: a multicentre randomised trial - ClinicalKey (no date). Available at: <https://www-clinicalkey-com.ezproxy4.lib.le.ac.uk/#!/content/journal/1-s2.0-S0140673610606320>.

'Professor Kamlesh Khunti - Coding, Classification and Diagnosis of Diabetes' (4AD). Available at: <https://www.youtube.com/watch?v=AhhWTmEFuag>.

Raccah, D., Huet, D., Dib, A, et al. (2017a) 'Review of basal-plus insulin regimen options for simpler insulin intensification in people with Type 2 diabetes mellitus', *Diabetic Medicine* [Preprint]. Available at: <https://doi.org/10.1111/dme.13390>.

- Raccah, D., Huet, D., Dib, A., et al. (2017b) 'Review of basal-plus insulin regimen options for simpler insulin intensification in people with Type 2 diabetes mellitus', *Diabetic Medicine* [Preprint]. Available at: <https://doi.org/10.1111/dme.13390>.
- Raccah, D., Huet, D., Dib, A., et al. (2017a) 'Review of basal-plus insulin regimen options for simpler insulin intensification in people with Type 2 diabetes mellitus', *Diabetic Medicine*, 34(9), pp. 1193–1204. Available at: <https://doi.org/10.1111/dme.13390>.
- Raccah, D., Huet, D., Dib, A., et al. (2017b) 'Review of basal-plus insulin regimen options for simpler insulin intensification in people with Type 2 diabetes mellitus', *Diabetic Medicine*, 34(9), pp. 1193–1204. Available at: <https://doi.org/10.1111/dme.13390>.
- Randomized controlled trial of insulin detemir versus NPH for the treatment of pregnant women with diabetes - ClinicalKey (no date). Available at: <https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S0002937815005931>.
- Richter, B. and Neises, G. (1996) "'Human" insulin versus animal insulin in people with diabetes mellitus', in *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd. Available at: <https://doi.org/10.1002/14651858.CD003816.pub2>.
- Riddle, M.C. et al. (2013) 'Adding Once-Daily Lixisenatide for Type 2 Diabetes Inadequately Controlled by Established Basal Insulin: A 24-week, randomized, placebo-controlled comparison (GetGoal-L)', *Diabetes Care*, 36(9), pp. 2489–2496. Available at: <https://doi.org/10.2337/dc12-2454>.
- Riddle, Matthew C. et al. (2014) 'New Insulin Glargine 300 Units/mL Versus Glargine 100 Units/mL in People With Type 2 Diabetes Using Basal and Mealtime Insulin: Glucose Control and Hypoglycemia in a 6-Month Randomized Controlled Trial (EDITION 1)', *Diabetes Care*, 37(10), pp. 2755–2762. Available at: <https://doi.org/10.2337/dc14-0991>.
- Riddle, M. C. et al. (2014) 'Randomized, 1-year comparison of three ways to initiate and advance insulin for type 2 diabetes: twice-daily premixed insulin versus basal insulin with either basal-plus one prandial insulin or basal-bolus up to three prandial injections', *Diabetes, Obesity and Metabolism*, 16(5), pp. 396–402. Available at: <https://doi.org/10.1111/dom.12225>.
- Rodbard, H.W. et al. (2017) 'Safety and efficacy of insulin degludec/liraglutide (IDegLira) added to sulphonylurea alone or to sulphonylurea and metformin in insulin-naïve people with Type 2 diabetes: the DUAL IV trial', *Diabetic Medicine*, 34(2), pp. 189–196. Available at: <https://doi.org/10.1111/dme.13256>.
- Rorsman, P. and Renström, E. (2003) 'Insulin granule dynamics in pancreatic beta cells', *Diabetologia*, 46(8), pp. 1029–1045. Available at: <https://doi.org/10.1007/s00125-003-1153-1>.
- Rosenstock, J., Bergenstal, R.M., et al. (2013) 'Better Glycemic Control and Weight Loss With the Novel Long-Acting Basal Insulin LY2605541 Compared With Insulin Glargine in Type 1 Diabetes: A randomized, crossover study', *Diabetes Care*, 36(3), pp. 522–528. Available at: <https://doi.org/10.2337/dc12-0067>.
- Rosenstock, J., Raccah, D., et al. (2013) 'Efficacy and Safety of Lixisenatide Once Daily

Versus Exenatide Twice Daily in Type 2 Diabetes Inadequately Controlled on Metformin: A 24-week, randomized, open-label, active-controlled study (GetGoal-X)', *Diabetes Care*, 36(10), pp. 2945–2951. Available at: <https://doi.org/10.2337/dc12-2709>.

Rosenstock, J. et al. (2014a) 'Improved Glucose Control With Weight Loss, Lower Insulin Doses, and No Increased Hypoglycemia With Empagliflozin Added to Titrated Multiple Daily Injections of Insulin in Obese Inadequately Controlled Type 2 Diabetes', *Diabetes Care*, 37(7), pp. 1815–1823. Available at: <https://doi.org/10.2337/dc13-3055>.

Rosenstock, J. et al. (2014b) 'Improved Glucose Control With Weight Loss, Lower Insulin Doses, and No Increased Hypoglycemia With Empagliflozin Added to Titrated Multiple Daily Injections of Insulin in Obese Inadequately Controlled Type 2 Diabetes', *Diabetes Care*, 37(7), pp. 1815–1823. Available at: <https://doi.org/10.2337/dc13-3055>.

Rowles, S., Kilvert, A. and Sinclair, A. (2011a) 'ABCD position statement on diabetes and end of life care', *Practical Diabetes International*, 28(1), pp. 26–27. Available at: <https://doi.org/10.1002/pdi.1547>.

Rowles, S., Kilvert, A. and Sinclair, A. (2011b) 'ABCD position statement on diabetes and end of life care', *Practical Diabetes International*, 28(1), pp. 26–27. Available at: <https://doi.org/10.1002/pdi.1547>.

Russell-Jones, D. and Khan, R. (2007) 'Insulin-associated weight gain in diabetes – causes, effects and coping strategies', *Diabetes, Obesity and Metabolism*, 9(6), pp. 799–812. Available at: <https://doi.org/10.1111/j.1463-1326.2006.00686.x>.

Shosuke Satake, Mary Courtney Moore, Kayano Igawa, Margaret Converse, Benjamin Farmer, Doss W. Neal (no date) 'Direct and indirect effects of insulin on glucose uptake and storage by the liver. (Original Articles)', *Diabetes [Preprint]*. Available at: http://go.galegroup.com.ezproxy4.lib.le.ac.uk/ps/retrieve.do?tabID=T002&resultListType=RESULT_LIST&searchResultsType=SingleTab&searchType=AdvancedSearchForm¤tPosition=2&docId=GALE%7CA87347789&docType=Article&sort=Relevance&contentSegment=&prodId=EAIM&contentSet=GALE%7CA87347789&searchId=R8&userGroupName=leicester&inPS=true.

Simmons, J.H. et al. (2007) 'Reliability of the Diabetes Fear of Injecting and Self-Testing Questionnaire in Pediatric Patients With Type 1 Diabetes', *Diabetes Care*, 30(4), pp. 987–988. Available at: <https://doi.org/10.2337/dc06-1553>.

Simmons, L.R. et al. (2012) 'Steroid-Induced Diabetes: Is It Just Unmasking of Type 2 Diabetes?', *ISRN Endocrinology*, 2012, pp. 1–5. Available at: <https://doi.org/10.5402/2012/910905>.

Srinivasan, B.T. et al. (2008) 'Recent advances in the management of type 2 diabetes mellitus: a review', *Postgraduate Medical Journal*, 84(996), pp. 524–531. Available at: <https://doi.org/10.1136/pgmj.2008.067918>.

Srinivasan, B.T. and Davies, M. (2014) 'Glycaemic management of type 2 diabetes', *Medicine*, 42(12), pp. 711–717. Available at: <https://doi.org/10.1016/j.mpmed.2014.09.011>.

Suggested insulin regimens for patients with type 1 diabetes mellitus who wish to fast

during the month of Ramadan - ClinicalKey (no date). Available at: <https://www-clinicalkey-com.ezproxy3.lib.le.ac.uk/#!/content/journal/1-s2.0-S0149291808002737>.

Swinnen, S.G. et al. (1996) 'Insulin detemir versus insulin glargine for type 2 diabetes mellitus', in Cochrane Database of Systematic Reviews. Chichester, UK: John Wiley & Sons, Ltd. Available at: <https://doi.org/10.1002/14651858.CD006383.pub2>.

Tapley, M. and Needham, E. (2012) 'Improving end of life care for people with diabetes', *Practical Diabetes*, 29(8), pp. 306–307a. Available at: <https://doi.org/10.1002/pdi.1711>.

'The Effect of Intensive Treatment of Diabetes on the Development and Progression of Long-Term Complications in Insulin-Dependent Diabetes Mellitus' (1993) *New England Journal of Medicine*, 329(14), pp. 977–986. Available at: <https://doi.org/10.1056/NEJM199309303291401>.

Tikkanen-Dolenc, H. et al. (2016) 'Frequent and intensive physical activity reduces risk of cardiovascular events in type 1 diabetes', *Diabetologia* [Preprint]. Available at: <https://doi.org/10.1007/s00125-016-4189-8>.

'Training in flexible, intensive insulin management to enable dietary freedom in people with type 1 diabetes: dose adjustment for normal eating (DAFNE) randomised controlled trial' (2002) *BMJ*, 325(7367), pp. 746–746. Available at: <https://doi.org/10.1136/bmj.325.7367.746>.

Villani, M., de Courten, B. and Zoungas, S. (2017) 'Emergency treatment of hypoglycaemia: a guideline and evidence review', *Diabetic Medicine*, 34(9), pp. 1205–1211. Available at: <https://doi.org/10.1111/dme.13379>.

W. Blair Geho (2014) 'A Single-blind, Placebo-controlled, Dose-ranging Trial of Oral Hepatic-directed Vesicle Insulin Add-on to Oral Antidiabetic Treatment in Patients With Type 2 Diabetes Mellitus', *Journal of Diabetes Science and Technology*, 8(3). Available at: <https://doi.org/doi:10.1177/1932296814524871>.

Wahren, J. and Kallas, A. (2012) 'Loss of Pulsatile Insulin Secretion: A Factor in the Pathogenesis of Type 2 Diabetes?', *Diabetes*, 61(9), pp. 2228–2229. Available at: <https://doi.org/10.2337/db12-0664>.

Walker, R.A., Rodgers, J., and Diabetes UK. (2010) *Diabetes: a practical guide to managing your health*. Fully revised and updated. London: Dorling Kindersley.

Wang, C., Mamza, J. and Idris, I. (2015) 'Biphasic vs basal bolus insulin regimen in Type 2 diabetes: a systematic review and meta-analysis of randomized controlled trials', *Diabetic Medicine*, 32(5), pp. 585–594. Available at: <https://doi.org/10.1111/dme.12694>.

Wang, W. et al. (2017) 'Effects of Insulin Plus Glucagon-Like Peptide-1 Receptor Agonists (GLP-1RAs) in Treating Type 1 Diabetes Mellitus: A Systematic Review and Meta-Analysis', *Diabetes Therapy* [Preprint]. Available at: <https://doi.org/10.1007/s13300-017-0282-3>.

Yki-Jarvinen, H. et al. (2007) 'Initiate Insulin by Aggressive Titration and Education (INITIATE): A randomized study to compare initiation of insulin combination therapy in type 2 diabetic patients individually and in groups', *Diabetes Care*, 30(6), pp. 1364–1369.

Available at: <https://doi.org/10.2337/dc06-1357>.

Yki-Jarvinen, H. and Kotronen, A. (2013) 'Is There Evidence to Support Use of Premixed or Prandial Insulin Regimens in Insulin-Naive or Previously Insulin-Treated Type 2 Diabetic Patients?', *Diabetes Care*, 36(Supplement_2), pp. S205–S211. Available at: <https://doi.org/10.2337/dcS13-2026>.

Young, L.A. and Buse, J.B. (2014) 'GLP-1 receptor agonists and basal insulin in type 2 diabetes', *The Lancet*, 384(9961), pp. 2180–2181. Available at: [https://doi.org/10.1016/S0140-6736\(14\)61409-4](https://doi.org/10.1016/S0140-6736(14)61409-4).

Zambanini, A. et al. (1999) 'Injection related anxiety in insulin-treated diabetes', *Diabetes Research and Clinical Practice*, 46(3), pp. 239–246. Available at: [https://doi.org/10.1016/S0168-8227\(99\)00099-6](https://doi.org/10.1016/S0168-8227(99)00099-6).

Zinman, B. et al. (2011) 'Insulin degludec, an ultra-long-acting basal insulin, once a day or three times a week versus insulin glargine once a day in patients with type 2 diabetes: a 16-week, randomised, open-label, phase 2 trial', *The Lancet*, 377(9769), pp. 924–931. Available at: [https://doi.org/10.1016/S0140-6736\(10\)62305-7](https://doi.org/10.1016/S0140-6736(10)62305-7).